APPROXIMATION OF ABSTRACT DIFFERENTIAL EQUATIONS

Davide Guidetti, Bülent Karasözen and Serguei Piskarev

April 25, 2002

Abstract

This review paper is devoted to the numerical analysis of abstract differential equations in Banach spaces. The presentation is given on general approximation scheme and based on semigroup theory and functional analysis approach. The paper concerns mainly numerical analysis of differential equations of first order in time.

1991 Mathematics Subject Classification: 65J, 65M

Keywords and phrases: Abstract differential equations, C_0-semigroups, Banach spaces, semidiscretization, coerciveness, maximal regularity inequality, Lax's equivalence theorem, difference schemes, semilinear equations, approximation of the spectrum of operators, discrete semigroups, rational approximations, backward evolution equations, ill-posed problems, stochastic regularization

Contents

1 Introduction **2**

2 General approximation scheme **2**

2.1 Approximation of spectrum of linear operators 3
2.2 Regions of convergence 4
2.3 Convergence in one space dimension 5
2.4 Compact convergence of resolvents 6

3 Discretization of semigroups **7**

3.1 The simplest discretization schemes 7
3.2 Rational approximation 10
3.3 Richardson's Extrapolation method 12
3.4 Lax-type equivalence theorems with orders 12

4 Backward Cauchy problem **15**

4.1 C-semigroups and ill-posed problems 16
4.2 Semidiscrete approximation theorem 18
4.3 Approximation by discrete C-semigroups 19

5 Coercive inequalities **21**
5.1 Coercive inequality in $C_{\tau_n}([0,T];E_n)$ spaces 22
5.2 Coercive inequality in $C^{\alpha,0}_{\tau_n}([0,T];E_n)$ spaces 23
5.3 Coercive inequality in $L^p_{\tau_n}([0,T];E_n)$ spaces 23
5.4 Coercive inequality in $B_{\tau_n}\left([0,T];C^\theta_h(\Omega_h)\right) \cap C_h([0,T];C_h(\bar{\Omega}_h))$ 26

6 Approximations of Semilinear Equations **31**
6.1 Approximations of Cauchy problem 31
6.2 Approximation of periodic problem 32

1 Introduction

This review paper is devoted to the numerical analysis of abstract differential equations in Banach spaces. Most of the finite difference, finite element and projection methods could be considered from the point of view of general approximation schemes (see for example [204], [208], [207] for such a representation). Results obtained for general approximation schemes make the formulation of concrete numerical methods easier and gives an overview of methods which are suitable for different classes of problems.

The qualitative theory of differential equations in Banach spaces presented in many brilliant papers and books. We can refer to the bibliography [216] which contains about 2500 references. Unfortunately there are no books or reviews on approximation theory for differential equations in abstract spaces last 20 years. Any information on the subject can be found only in the original papers. It looks that such a review is the first step on the way to describe a complete picture of discretisation methods for abstract differential equations in Banach spaces.

In Section 2 we describe the general approximation scheme, different types of convergence of operators and the relation of convergence with the approximation of spectra. Also such a convergence analysis could be used if one considers elliptic problems, i.e. the problems which do not depend on time.

Section 3 contains a complete picture of the theory of discretization of semigroups in Banach spaces. It summarizes Trotter-Kato and Lax-Richtmyer theorems from the general and common point of view and related problems.

Approximation of ill-posed problems is considered in Section 4, which are based on the theory of approximation of local C-semigroups. Since the backward Cauchy problem is very important in applications and admits stochastic noise, we consider also approximation using stochastic regularization. Such an approach has never been considered in the literature before to the best our knowledge.

In Section 5 we present discrete coercive inequalities for abstract parabolic equations in $C_{\tau_n}([0,T];E_n)$, $C^\alpha_{\tau_n}([0,T];E_n)$, $L^p_{\tau_n}([0,T];E_n)$ and $B_{\tau_n}([0,T];C^\alpha(\Omega_h))$ spaces.

The last Section 6 deals with semilinear problems. We consider approximations of Cauchy problems and also the problems with periodic solutions. The approach we describe is based on the theory of rotation of vector fields and the principle of compact approximation of operators.

2 General approximation scheme

Let $B(E)$ denote the Banach algebra of all linear bounded operators on a complex Banach space E. The set of all linear closed densely defined operators in E will be denoted by $\mathcal{C}(E)$. We denote by $\sigma(B)$ the spectrum of the operator B, by $\rho(B)$ the resolvent set of B, by $\mathcal{N}(B)$ the null space of B and by $\mathcal{R}(B)$ the range of B. Let us recall that $B \in B(E)$ is said to be a Fredholm operator if $\mathcal{R}(B)$ is closed, $dim\,\mathcal{N}(B) < \infty$ and $codim\,\mathcal{R}(B) < \infty$, the index of B is defined as $ind\,B = dim\,\mathcal{N}(B) - codim\,\mathcal{R}(B)$.

The general approximation scheme, due to [80]-[82], [184], [207], [204] can be described in the following way. Let E_n and E be Banach spaces and $\{p_n\}$ be a sequence of linear bounded operators $p_n : E \to E_n, p_n \in B(E, E_n), n \in I\!N = \{1, 2, \cdots\}$, with the property:

$$\|p_n x\|_{E_n} \to \|x\|_E \text{ as } n \to \infty \text{ for any } x \in E.$$

Definition 2.1 *The sequence of elements $\{x_n\}, x_n \in E_n, n \in I\!N$, is said to be $\mathcal{P}$-convergent to $x \in E$ iff $\|x_n - p_n x\|_{E_n} \to 0$ as $n \to \infty$ and we write this $x_n \to x$.*

Definition 2.2 *The sequence of elements $\{x_n\}, x_n \in E_n, n \in I\!N$, is said to be $\mathcal{P}$-compact if for any $I\!N' \subseteq I\!N$ there exist $I\!N'' \subseteq I\!N'$ and $x \in E$ such that $x_n \to x$, as $n \to \infty$ in $I\!N''$.*

Definition 2.3 *The sequence of bounded linear operators $B_n \in B(E_n), n \in I\!N$, is said to be $\mathcal{PP}$-convergent to the bounded linear operator $B \in B(E)$ if for every $x \in E$ and for every sequence $\{x_n\}, x_n \in E_n, n \in I\!N$, such that $x_n \to x$ one has $B_n x_n \to Bx$. We write then $B_n \to B$.*

For general examples of notions of $\mathcal{P}-$convergence see [79], [184], [200], [208].

Remark 2.1 *If we put $E_n = E$ and $p_n = I$ for each $n \in I\!N$, where I is the identity operator on E, then Definition 2.1 leads to the traditional pointwise convergent bounded linear operators which we denote by $B_n \to B$.*

Let us denote by E^+ the positive cone in a Banach lattice E. The operator B is called positive if for any $x^+ \in E^+$ it follows $Bx^+ \in E^+$; we write $0 \preceq B$.

Definition 2.4 *The system $\{p_n\}$ is called discrete order preserving if for all sequences $\{x_n\}, x_n \in E_n$, and element $x \in E$ the following implication holds: $x_n \to x$ implies $x_n^+ \to x^+$.*

It is known [96] that $\{p_n\}$ preserves the order iff $\|p_n x^+ - (p_n x)^+\|_{E_n} \to 0$ as $n \to \infty$ for any $x \in E$.

In the case of unbounded operators, and we know in general infinitesimal generators are unbounded, we consider the notion of *compatibility*.

Definition 2.5 *The sequence of closed linear operators $\{A_n\}, A_n \in \mathcal{C}(E_n), n \in I\!N$, are said to be compatible with a closed linear operator $A \in \mathcal{C}(E)$ iff for each $x \in D(A)$ there is a sequence $\{x_n\}, x_n \in D(A_n) \subseteq E_n, n \in I\!N$, such that $x_n \to x$ and $A_n x_n \to Ax$. We write (A_n, A) are compatible.*

Usually in practice Banach spaces E_n are finite dimensional, although, in general, say for the case of a closed operator A, we have $\dim E_n \to \infty$ and $\|A_n\|_{B(E_n)} \to \infty$ as $n \to \infty$.

2.1 Approximation of spectrum of linear operators

The most important role in approximations of equation $Bx = y$ and approximations of spectra of operator B, is played by the notions of *stable* and *regular* convergence. These notions are used in different areas of numerical analysis (cf. [10], [15], [78], [83]-[86], [207], [209], [204], [220]).

Definition 2.6 *A sequence of operators $\{B_n\}, B_n \in B(E_n), n \in I\!N$, is stably convergence to an operator $B \in B(E)$ iff $B_n \to B$ and $\|B_n^{-1}\|_{B(E_n)} = O(1), n \to \infty$. We will write $B_n \to B$ stably.*

Definition 2.7 *A sequence of operators $\{B_n\}, B_n \in B(E_n)$, is called regularly convergent to the operator $B \in B(E)$ iff $B_n \to B$ and the following implication holds:*

$$\|x_n\|_{E_n} = O(1) \ \& \ \{B_n x_n\} \text{ is } \mathcal{P} - \text{compact} \Longrightarrow \{x_n\} \text{ is } \mathcal{P}\text{-compact.}$$

We write this as $B_n \to B$ regularly.

Theorem 2.1 *[207] For $B_n \in B(E_n)$ and $B \in B(E)$ the following conditions are equivalent:*

(i) $B_n \to B$ regularly, B_n are Fredholm operators of index 0 and $\mathcal{N}(B) = \{0\}$;

(ii) $B_n \to B$ stably and $\mathcal{R}(B) = E$;

(iii) $B_n \to B$ stably and regularly;

(iv) if one of the conditions (i)-(iii) is fulfilled, then there exist $B_n^{-1} \in B(E_n), B^{-1} \in B(E)$ and $B_n^{-1} \to B^{-1}$ regularly and stably.

This theorem admits an extension to the case of closed operators $B \in \mathcal{C}(E), B_n \in \mathcal{C}(E_n)$ [210].

Let $\Lambda \subseteq \mathbb{C}$ be some open connected set and let $B \in B(E)$. For an isolated point $\lambda \in \sigma(B)$, the corresponding maximal invariant space (or generalized eigenspace) will be denoted by $\mathcal{W}(\lambda; B) = P(\lambda)E$, where $P(\lambda) = \frac{1}{2\pi i}\int_{|\zeta-\lambda|=\delta}(\zeta I - B)^{-1}d\zeta$ and δ is small enough so that there are no points of $\sigma(B)$ in the disc $\{\zeta : |\zeta - \lambda| \leq \delta\}$ other than λ. The isolated point $\lambda \in \sigma(B)$ is a *Riesz point* of B if $\lambda I - B$ is a Fredholm operator with index zero and $P(\lambda)$ is of finite rank. Denote by $\mathcal{W}(\lambda, \delta; B_n) = \bigcup_{|\lambda_n - \lambda| < \delta, \lambda_n \in \sigma(B_n)} \mathcal{W}(\lambda_n, B_n)$, where $\lambda_n \in \sigma(B_n)$ are taken from a δ-neighborhood of λ. It is clear that $\mathcal{W}(\lambda, \delta; B_n) = P_n(\lambda)E_n$, where $P_n(\lambda) = \frac{1}{2\pi i}\int_{|\zeta-\lambda|=\delta}(\zeta I - B_n)^{-1}d\zeta$. The following theorems state the complete picture of the approximation of the spectrum.

Theorem 2.2 *[79], [205]-[206] Assume that $L_n(\lambda) = \lambda I - B_n$ and $L(\lambda) = \lambda I - B$ are Fredholm operators with index zero for any $\lambda \in \Lambda$ and that $L_n(\lambda) \to L(\lambda)$ stably for any $\lambda \in \rho(B) \cap \Lambda \neq \emptyset$. Then*

(i) for any $\lambda_0 \in \sigma(B) \cap \Lambda$ there exists a sequence $\{\lambda_n\}$, $\lambda_n \in \sigma(B_n)$, $n \in \mathbb{N}$, such that $\lambda_n \to \lambda_0$ as $n \to \infty$;

(ii) if for some sequence $\{\lambda_n\}, \lambda_n \in \sigma(B_n)$, $n \in \mathbb{N}$, one has $\lambda_n \to \lambda_0 \in \Lambda$ as $n \to \infty$, then $\lambda_0 \in \sigma(B)$;

(iii) for any $x \in \mathcal{W}(\lambda_0, B)$ there exists a sequence $\{x_n\}$, $x_n \in \mathcal{W}(\lambda_0, \delta; B_n)$, $n \in \mathbb{N}$, such that $x_n \to x$ as $n \to \infty$;

(iv) there exists $n_0 \in \mathbb{N}$ such that $\dim \mathcal{W}(\lambda_0, \delta; B_n) \geq \dim \mathcal{W}(\lambda_0, B)$ for any $n \geq n_0$.

Remark 2.2 *The inequality in (iv) can be strict for all $n \in \mathbb{N}$ as is shown in [204].*

Theorem 2.3 *[207] Assume that $L_n(\lambda)$ and $L(\lambda)$ are Fredholm operators with index zero for all $\lambda \in \Lambda$. Suppose that $L_n(\lambda) \to L(\lambda)$ regularly for any $\lambda \in \Lambda$ and that $\rho(B) \cap \Lambda \neq \emptyset$. Then the statements (i)-(iii) of Theorem 2.2 hold and also*

(iv) there exists $n_0 \in \mathbb{N}$ such that $\dim \mathcal{W}(\lambda_0, \delta; B_n) = \dim \mathcal{W}(\lambda_0, B)$ for all $n \geq n_0$;

(v) any sequence $\{x_n\}, x_n \in \mathcal{W}(\lambda_0, \delta; B_n)$, $n \in \mathbb{N}$, with $\|x_n\|_{E_n} = 1$ is $\mathcal{P}$-compact and any limit point of this sequence belongs to $\mathcal{W}(\lambda_0, B)$.

Remark 2.3 *Estimates of $|\lambda_n - \lambda_0|$, $\operatorname{gap}\left(\mathcal{W}(\lambda_0, \delta; B_n), \mathcal{W}(\lambda_0, B)\right)$ and $|\hat{\lambda}_n - \lambda_0|$ are given in [207], where $\hat{\lambda}_n$ denotes the arithmetic mean (counting algebraic multiplicities) of the spectral values of B_n that contribute to $\mathcal{W}(\lambda_0, \delta; B_n)$. For the notion of gap and its properties, see [102].*

2.2 Regions of convergence

Theorems 2.2 and 2.3 have been generalized in [210] to the case of closed operators using the following notions introduced by Kato [102].

Definition 2.8 *The region of stability $\Delta_s = \Delta_s(\{A_n\})$, $A_n \in \mathcal{C}(B_n)$, is defined to be the set of all $\lambda \in \mathbb{C}$ such that $\lambda \in \rho(A_n)$ for almost all n and such that the sequence $\{\|(\lambda I - A_n)^{-1}\|\}_{n \in \mathbb{N}}$ is bounded. The region of convergence $\Delta_c = \Delta_c(\{A_n\})$, $A_n \in \mathcal{C}(E_n)$, is defined as the set of all $\lambda \in \mathbb{C}$ such that $\lambda \in \Delta_s(\{A_n\})$ and such that the sequence of operators $\{(\lambda I - A_n)^{-1}\}_{n \in \mathbb{N}}$ is $\mathcal{PP}$-convergent to some operator $S(\lambda) \in B(E)$.*

It is clear that $S(\cdot)$ is pseudo-resolvent and $S(\cdot)$ is a resolvent of some operator iff $\mathcal{N}(S(\lambda)) = \{0\}$ for some λ (cf. [102]).

Definition 2.9 *A sequence of operators $\{K_n\}$, $K_n \in \mathcal{C}(E_n)$, is called regularly compatible with the operator $K \in \mathcal{C}(E)$ if (K_n, K) are compatible and, for any bounded sequence $\|x_n\|_{E_n} = O(1)$ such that $x_n \in D(K_n)$ and such that $\{K_n x_n\}$ is $\mathcal{P}$-compact, it follows that $\{x_n\}$ is $\mathcal{P}$-compact and the $\mathcal{P}$-convergence of $\{x_n\}$ to some x and of $\{K_n x_n\}$ to some y for $n \to \infty$ in $I\!N' \subseteq I\!N$ imply that $x \in D(K)$ and $Kx = y$.*

Definition 2.10 *The region of regularity $\Delta_r = \Delta_r(\{A_n\}, A)$, is defined as the set of all $\lambda \in \mathbb{C}$ such that (K_n, K) are regularly compatible, where $K_n = \lambda I - A_n$ and $K = \lambda I - A$.*

The relationships between these regions are given by

Proposition 2.1 *[210] Suppose that $\Delta_c \neq \emptyset$ and $\mathcal{N}\big(S(\lambda)\big) = \{0\}$ at least for one point $\lambda \in \Delta_c$ so that $S(\lambda) = (I\lambda - A)^{-1}$. Then (A_n, A) are compatible and*

$$\Delta_c = \Delta_s \cap \rho(A) = \Delta_s \cap \Delta_r = \Delta_r \cap \rho(A).$$

It is shown in [210] that the following three conditions : *(A_n, A) are compatible, $\lambda I - A_n$ and $\lambda I - A$ are Fredholm operators with index zero for any $\lambda \in \Lambda$ and $\rho(A) \cap \Lambda \neq \emptyset$* imply (i)-(iv) of Theorem 2.2 when $\rho(A) \cap \Lambda \subseteq \Delta_s$ and imply (i)-(iii) of Theorem 2.2 and (iv)-(v) of Theorem 2.3, when $\Lambda \subseteq \Delta_r$.

Definition 2.11 *The Riesz point $\lambda_0 \in \sigma(A)$ is said to be strongly stable in Kato's sense if* $\dim \mathcal{W}(\lambda_0, \delta; B_n) \leq \dim \mathcal{W}(\lambda_0, B)$ *for all $n \geq n_0$.*

Theorem 2.4 *[210] The Riesz point $\lambda_0 \in \sigma(A)$ is strongly stable in Kato's sense iff $\lambda_0 \in \Lambda \cap \Delta_r \cap \sigma(A)$.*

Investigations of approximation of spectra and types of convergence, but not on general approximation scheme are given in [13], [37], [49], [127], [128], [139], [142].

2.3 Convergence in one space dimension

Throughout this subsection we assume that $E_n = E$ and $p_n = I$ for all $n \in I\!N$. Hence, the symbol $\mathcal{P}$ will be omitted in all notations of this subsection.

Let us recall that if $B_n \to B$ compactly (see Definition 2.12), then for any $\lambda \neq 0$ we have $\lambda I - B_n \to \lambda I - B$ regularly [204]. When $B_n \to B$ compactly and B is a compact operator, Anselone [10] has proved that

$$\|(B_n - B)B_n\| \to 0, \ \|(B_n - B)B\| \to 0 \ \text{ as } n \to \infty. \tag{2.1}$$

Considering an approximation of a weakly singular compact integral operator Ahues [4] has proved that these convergence properties (2.1) are sufficient to state that a Riesz point is strongly stable in Kato's sense.

Theorem 2.5 *[7] Assume that $B \in B(E)$ is compact and that $B_n \to B$. If $\|(B_n - B)B_n\| \to 0$ as $n \to \infty$ then for any nonzero $\lambda_0 \in \sigma(B)$ the assertions (i)-(iii) of Theorem 2.2 and assertions (iv)-(v) of Theorem 2.3 hold.*

Theorem 2.6 *[7] Assume that $B_n \to B$ and (2.1) holds. Then for any non-zero Riesz point $\lambda_0 \in \sigma(B)$ the assertions (i)-(iii) of Theorem 2.2 and assertions (iv)-(v) of Theorem 2.5 hold.*

Corollary 2.1 *[5] Assume that $B_n \to B$, that $\lambda I - B_n$ are Fredholm operators with index zero for $\lambda \in \{z : |z-\lambda_0| \le \delta\}$ and $\|(B_n - B)B_n^k\| \to 0$ as $n \to \infty$ for some $k \in I\!N$. Then for any non zero Riesz point $\lambda_0 \in \sigma(B)$ the assertions (i)-(iii) of Theorem 2.2 and assertions (iv)-(v) of Theorem 2.3 hold.*

Theorem 2.7 *[7] Assume that B is compact, $B_n \to B$ and $\|B_n(B_n - B)\| \to 0$. Then $\lambda_0 I - B_n \to \lambda_0 I - B$ regularly for any $\lambda_0 \neq 0$.*

Theorem 2.8 *[7] Assume that B is compact, $B_n \to B$ and $\|B_n^k(B_n - B)\| \to 0$ for some $k \in I\!N$. Then $\lambda_0 I - B_n \to \lambda_0 I - B$ regularly for any $\lambda_0 \neq 0$.*

Let $r(B)$ be a spectral radius of operator $B \in B(E)$.

Theorem 2.9 *[18] Let E be a Banach lattice. Let $0 \preceq B_n, B \in B(E)$ be such that $B_n \to B$ and $\|(B_n - B)^+\| \to 0$ as $n \to \infty$. Suppose that $r(B)$ is a Riesz point of $\sigma(B)$. Then $r(B_n)$ is a Riesz point of $\sigma(B_n)$ and $r(B_n) \to r(B)$ as $n \to \infty$.*

The conclusion on order of convergence of eigenvectors in Theorem 2.9 also is given in [17].

The application of Theorems 2.5 - 2.8 to the numerical solution of a mathematical model used in the jet printer industry is considered [6], [115].

2.4 Compact convergence of resolvents

We consider now the important class of operators which have compact resolvents. We will use this property of generator as assumption in section 6. It is natural to consider in this case approximate operators which "keep" this property.

Definition 2.12 *A sequence of operators $\{B_n\}$, $B_n \in B(E_n)$, $n \in I\!N$, converges compactly to an operator $B \in B(E)$, if $B_n \to B$ and the following compactness condition holds:*

$$\|x_n\|_{E_n} = O(1) \Longrightarrow \{B_n x_n\} \text{ is } \mathcal{P}\text{-compact.}$$

Definition 2.13 *The region of compact convergence of resolvents, $\Delta_{cc} = \Delta_{cc}(A_n, A)$, where $A_n \in \mathcal{C}(E_n)$ and $A \in \mathcal{C}(E)$ is defined as the set of all $\lambda \in \Delta_c \cap \rho(A)$ such that $(\lambda I - A_n)^{-1} \to (\lambda I - A)^{-1}$ compactly.*

Theorem 2.10 *Assume that $\Delta_{cc} \neq \emptyset$. Then for any $\zeta \in \Delta_s$ the following implication holds:*

$$\|x_n\|_{E_n} = O(1) \ \& \ \|(\zeta I - A_n)x_n\|_{E_n} = O(1) \Longrightarrow \{x_n\} \text{ is } \mathcal{P}\text{-compact.} \tag{2.2}$$

Conversely, if for some $\zeta \in \Delta_c \cap \rho(A)$ the implication (2.2) holds, then $\Delta_{cc} \neq \emptyset$.

Proof. Let $(\mu I - A_n)^{-1} \to (\mu I - A)^{-1}$compactly for some $\mu \in \Delta_{cc}$. Then for $\|x_n\|_{E_n} = O(1)$ and $\|(\zeta I - A_n)x_n\|_{E_n} = O(1)$, we get from Hilbert's identity

$$(\zeta I - A_n)^{-1} - (\mu I - A_n)^{-1} = (\mu - \zeta)(\zeta I - A_n)^{-1}(\mu I - A_n)^{-1}, \tag{2.3}$$

that $x_n = (\mu I - A_n)^{-1}(\zeta I - A_n)x_n - (\zeta - \mu)(\mu I - A_n)^{-1}x_n$ and it follows that $\{x_n\}$ is $\mathcal{P}$-compact. Conversely, let the implication (2.2) hold for some $\zeta_0 \in \Delta_c \cap \rho(A)$. We are going to show then that $\zeta_0 \in \Delta_{cc}$. Taking a bounded sequence $\{y_n\}$, $n \in I\!N$, one gets that the sequence $\|(\zeta_0 I - A_n)^{-1}y_n\|_{E_n} = O(1)$ for $n \in I\!N$. Let us apply the implication (2.2) to the sequence $x_n = (\zeta_0 I - A_n)^{-1}y_n$. It is easy to see that $\{x_n\}$ is $\mathcal{P}$-compact. Hence $\zeta_0 \in \Delta_{cc}$. □

Corollary 2.2 *Assume that* $\Delta_{cc} \neq \emptyset$. *Then* $\Delta_{cc} = \Delta_c \cap \rho(A)$.

Proof. It is clear that $\Delta_{cc} \subseteq \Delta_c \cap \rho(A)$. To prove that $\Delta_{cc} \supseteq \Delta_c \cap \rho(A)$ let us consider the Hilbert identity (2.3). Now let $\mu \in \Delta_{cc}$. Then $\mu \in \Delta_{cc} \cap \Delta_c \cap \rho(A)$. Hence for every $\zeta \in \Delta_c \cap \rho(A)$ and for any bounded sequence $\{x_n\}$, $n \in I\!N$, the sequence $\{(\zeta I - A_n)^{-1} x_n\}$ is $\mathcal{P}$-compact. □

Comparing Definitions 2.7, 2.8, 2.13 and implication (2.2) we see that $\Delta_{cc} \subseteq \Delta_r$.

Theorem 2.11 *Assume that* $\Delta_{cc} \neq \emptyset$. *Then* $\Delta_r = \mathbb{C}$.

Proof. Take any point $\lambda_1 \in \mathbb{C}$. We have to show that $(\lambda_1 I - A_n, \lambda_1 I - A)$ are regularly compatible. Assume that $\|x_n\|_{E_n} = O(1)$ and that $\{(\lambda_1 I - A_n)x_n\}$ is $\mathcal{P}$-compact. To show that $\{x_n\}$ is $\mathcal{P}$-compact take $\mu \in \Delta_{cc}$. Using (2.3) with $\zeta = \lambda_1$ one obtains $x_n = (\mu I - A_n)^{-1}(\lambda_1 I - A_n)x_n + (\lambda_1 - \mu)(\mu I - A_n)^{-1} x_n$ and therefore $\{x_n\}$ is $\mathcal{P}$-compact. Assume now that $x_n \to x$ and $(\lambda_1 I - A_n)^{-1} x_n \to y$, as $n \to \infty$ in $I\!N' \subseteq I\!N$. Then $x = (\mu I - A)^{-1} y - (\lambda_1 - \mu)(\mu I - A)^{-1} x$ and it follows that $x \in D(A)$ and $(\lambda_1 I - A)x = y$. □

3 Discretization of semigroups

Let us consider the well-posed Cauchy problem in the Banach space E with operator $A \in \mathcal{C}(E)$

$$\begin{aligned} u'(t) &= Au(t),\ t \in [0, \infty), \\ u(0) &= u^0, \end{aligned} \tag{3.1}$$

where operator A generates C_0-semigroup $\exp(\cdot A)$. It is well-known that the C_0-semigroup gives the solution of (3.1) by the formula $u(t) = \exp(tA)u^0$ for $t \geq 0$. The theory of well-posed problems and numerical analysis of these problems have been developed extensively, see for instance [72], [85], [102], [158], [160], [197], [213].

Let us consider on the general discretization scheme the semidiscrete approximation of the problem (3.1) in the Banach spaces E_n :

$$\begin{aligned} u_n'(t) &= A_n u_n(t),\ t \in [0, \infty), \\ u_n(0) &= u_n^0, \end{aligned} \quad . \tag{3.2}$$

with the operators $A_n \in \mathcal{C}(E_n)$, such that they generate C_0-semigroups, which are consistent with the operator A and $u_n^0 \to u^0$.

3.1 The simplest discretization schemes

We have the following version of Trotter-Kato's Theorem on general approximation scheme:

Theorem 3.1 *[200] (Theorem ABC) The following conditions (A) and (B) are equivalent to condition (C).*

(A) Consistency. There exists $\lambda \in \rho(A) \cap \cap_n \rho(A_n)$ *such that the resolvents converge*
$(\lambda I - A_n)^{-1} \to (\lambda - A)^{-1}$;

(B) Stability. There are some constants $M \geq 1$ *and* ω, *which are not depending on* n *and such that* $\|\exp(tA_n)\| \leq M \exp(\omega t)$ *for* $t \geq 0$ *and any* $n \in N$;

(C) Convergence. For any finite $T > 0$ *one has* $\max_{t \in [0,T]} \|\exp(tA_n)u_n^0 - p_n \exp(tA)u^0\| \to 0$ *as* $n \to \infty$, *whenever* $u_n^0 \to u^0$.

The analytic C_0-semigroup case is slightly different from the general case, but with the same property (A).

Theorem 3.2 *[158] Let operators A and A_n generate analytic C_0-semigroups. The following conditions (A) and (B_1) are equivalent to condition (C_1).*

(A) Consistency. There exists $\lambda \in \rho(A) \cap \cap_n \rho(A_n)$ such that the resolvents converge $(\lambda I - A_n)^{-1} \to (\lambda I - A)^{-1}$;

(B_1) Stability. There are some constants $M_2 \geq 1$ and ω_2 such that

$$\|(\lambda I - A_n)^{-1}\| \leq \frac{M_2}{|\lambda - \omega_2|}, \ Re\lambda > \omega_2, n \in N;$$

(C_1) Convergence. For any finite $\mu > 0$ and some $0 < \theta < \frac{\pi}{2}$ we have

$$\max_{\eta \in \Sigma(\theta,\mu)} \|\exp(\eta A_n)u_n^0 - p_n \exp(\eta A)u^0\| \to 0$$

as $n \to \infty$ whenever $u_n^0 \to u^0$. Here $\Sigma(\theta, \mu) = \{z \in \Sigma(\theta) : |z| \leq \mu\}$, and $\Sigma(\theta) = \{z \in \mathbb{C} \ : |arg\, z| \leq \theta\}$.

Definition 3.1 *The linear operator $A : D(A) \subseteq E \to E$ is said to have the positive off-diagonal (POD) property if $\langle Au, \phi\rangle \geq 0$ whenever $0 \preceq u \in D(A)$ and $0 \preceq \phi \in E^*$ with $\langle u, \phi\rangle = 0$.*

Definition 3.2 *An element $e \in E^+$ is said to be an order unit in E if for every $x \in E$ there exists $0 \leq \lambda \in R$ such that $-\lambda e \preceq x \preceq \lambda e$. For $e \in intE^+$ we can define an order unit norm by*

$$\|x\|_e = \inf\{\lambda \geq 0 : -\lambda e \preceq x \preceq \lambda e\}.$$

The ordered Banach space E is called an order unit space if there exists $e \in intE^+$ such that $\|\cdot\|_E = \|\cdot\|_e$.

We can state now Trotter-Kato's version for positive semigroups.

Theorem 3.3 *[166] Let the operators A_n and A from (3.1) and (3.2) be consistent and let E, E_n be order unit spaces and $e_n \in D(A_n) \cap intE_n^+$. Assume that the operators A_n have the POD property and $A_n e_n \preceq 0$ for sufficiently large n. Then $\exp(tA_n) \to \exp(tA)$ uniformly in $t \in [0, T]$.*

We can assume that conditions (A) and (B) for the corresponding semigroup case are satisfied without any restriction of generality if any discretization processes in time are considered. If we denote by $T_n(\cdot)$ a family of discrete semigroups as in [102], i.e. $\breve{A}_n = \frac{1}{\tau_n}(T_n(\tau_n) - I) \in B(E_n)$ and $T_n(t) = T_n(\tau_n)^{k_n}$, where $k_n = [\frac{t}{\tau_n}]$, as $\tau_n \to 0, n \to \infty$, then one gets

Theorem 3.4 *[200] (Theorem ABC-discr) The following conditions (A) and (B') are equivalent to condition (C').*

(A) Consistency. There exists $\lambda \in \rho(A) \cap \cap_n \rho(\breve{A}_n)$ such that the resolvents converge $(\lambda I - \breve{A}_n)^{-1} \to (\lambda I - A)^{-1}$;

(B') Stability. There are some constants $M_1 \geq 1$ and ω_1 such that

$$\|T_n(t)\| \leq M_1 \exp(\omega_1 t) \text{ for } t \in \overline{R}_+ = [0, \infty), n \in N;$$

(C') Convergence. For any finite $T > 0$ one has $\max_{t \in [0,T]} \|T_n(t)u_n^0 - p_n \exp(tA)u^0\| \to 0$ as $n \to \infty$, whenever $u_n^0 \to u^0$.

Theorem 3.5 *[200] Assume that conditions (A) and (B) of Theorem 3.1 hold. Then the implicit difference scheme*

$$\frac{\overline{U}_n(t+\tau_n)-\overline{U}_n(t)}{\tau_n}=A_n\overline{U}_n(t+\tau),\ \overline{U}_n(0)=u_n^0, \tag{3.3}$$

is stable, i.e. $\|(I_n-\tau_nA_n)^{-k_n}\|\le M_1e^{\omega_1 t}, t=k_n\tau_n\in\overline{IR}_+$, *and gives an approximation to the solution of the problem (3.1), i.e.* $\overline{U}_n(t)\equiv(I_n-\tau_nA_n)^{-k_n}u_n^0\to\exp(tA)u_n^0$ $\mathcal{P}$*-converges uniformly with respect to* $t=k_n\tau_n\in[0,T]$ *as* $u_n^0\to u^0, n\to\infty, k_n\to\infty, \tau_n\to 0$.

Here in Theorem 3.5 $\breve{A}_n=A_n(I_n-\tau_nA_n)^{-1}$ and therefore $(I_n-\tau_nA_n)^{-k_n}=(I_n+\tau_n\breve{A}_n)^{k_n}$.

Theorem 3.6 *[200] Assume that conditions (A) and (B) of the Theorem 3.1 hold and condition*

$$\tau_n\|A_n^2\|=O(1) \tag{3.4}$$

is fulfilled. Then the difference scheme

$$\frac{U_n(t+\tau_n)-U_n(t)}{\tau_n}=A_nU_n(t),\ U_n(0)=u_n^0, \tag{3.5}$$

is stable, i.e. $\|(I_n+\tau_nA_n)^{k_n}\|\le Me^{\omega t}, t=k_n\tau_n\in\overline{IR}_+$, *and gives an approximation to the solution of the problem (3.1), i.e.* $U_n(t)\equiv(I_n+\tau_nA_n)^{k_n}u_n^0\to u(t)$ $\mathcal{P}$*-converges uniformly with respect to* $t=k_n\tau_n\in[0,T]$ *as* $n\to\infty, k_n\to\infty, \tau_n\to 0$.

Theorem 3.7 *[158] Assume that conditions (A) and* (B_1) *of Theorem 3.2 hold and condition*

$$\tau_n\|A_n\|\le 1/(M+2), n\in I\!N \tag{3.6}$$

is fulfilled. Then the difference scheme (3.5) is stable and gives an approximation to the solution of the problem (3.1), i.e. $U_n(t)\equiv(I_n+\tau_nA_n)^{k_n}u_n^0\to u(t)$ *discretely* $\mathcal{P}$*-converge uniformly with respect to* $t=k_n\tau_n\in[0,T]$ *as* $u_n^0\to u^0, n\to\infty, k_n\to\infty, \tau_n\to 0$.

Let us introduce the following conditions:
(B_1') Stability. There are constants M',ω' such that

$$\|\exp(tA_n)\|\le M'e^{\omega' t},\ \|A_n\exp(tA_n)\|\le\frac{M'}{t}e^{\omega' t},\ t\in I\!R_+.$$

(B_1'') Stability. There are constants M'',ω'' and $\tau^*>0$ such that

$$\|(I-\tau_nA_n)^{-k}\|\le M''e^{\omega''k\tau_n},\ \|k\tau_nA_n(I-\tau_nA_n)^{-1}\|\le M''e^{\omega''k\tau_n},\ 0<\tau_n<\tau^*, n,k\in I\!N.$$

Proposition 3.1 *[180] Conditions* (B_1), (B_1'), (B_1'') *are equivalent.*

Theorem 3.8 *The conditions (A) and* (B_1'') *are equivalent to the condition* (C_1).

Theorem 3.9 *[161] Let the assumptions of Theorem 3.7 and (3.4) be satisfied. Then*

$$tA_n(I+\tau_nA_n)^{k_n}\to tA\exp(tA)\ \textit{uniformly in}\ t=k_n\tau_n\in[0,T]. \tag{3.7}$$

Conversely, if $(I+\tau_nA_n)^{k_n}\to\exp(tA)$ *uniformly in* $t=k_n\tau_n\in[0,T]$ *and (3.7) is satisfied, then condition* (C_1) *holds.*

Theorem 3.10 *[161] Let condition (B_1) hold. Then*

$$\|\exp(tA_n) - (I - \tau_n A_n)^{-k_n}\| \le c\frac{\tau_n}{t}e^{\omega t}.$$

If, moreover, stability condition (3.6) holds, then

$$\begin{aligned}
\|\exp(tA_n) - (I + \tau_n A_n)^{k_n}\| &\le c\frac{\tau_n}{t}e^{\omega t}, \\
\|(\exp(tA_n) - (I + \tau_n A_n)^{k_n})x_n\| &\le c\tau_n e^{\omega t}\|A_n x_n\|, \\
\|A_n(\exp(tA_n) - (I + \tau_n A_n)^{k_n})x_n\| &\le c\frac{\tau_n}{t}e^{\omega t}\|A_n x_n\|, \quad t = k_n\tau_n.
\end{aligned}$$

In the case of analytic C_0-semigroups for the forward scheme, as we saw, the following stability condition: $\tau_n\|A_n\| < 1/(M+2)$, which can't be improved even in Hilbert spaces for self-adjoint operators. In the case of almost periodic C_0-semigroups and the forward scheme for differential equations of first order in time (3.1), one obtains necessary and sufficient stability condition $\tau_n\|A_n\| < 1$ [160]. It was discovered that the stability condition of the forward scheme like (3.5) for the positive C_0-semigroups also can be written in the form $\tau_n\|A_n\| < 1$, see [165].

Stability of difference schemes for differential equations in Hilbert spaces in the energy norm are investigated in [176], [177], where also schemes with weights were considered. Semidiscrete approximations are studied in [177].

3.2 Rational approximation

Let us denote by $P_p(z)$ the element of the set of all real polynomials of degree not greater than p and by $\pi_{p,q}$ the set of all rational functions $r_{p,q}(z) = \frac{P_p(z)}{P_q(z)}$ and $P_q(0) = 1$. Then a Padé (p,q)-approximation for e^{-z} is defined as an element $R_{p,q}(z) \in \pi_{p,q}$ such that

$$|e^{-z} - R_{p,q}(z)| = O(|z|^{p+q+1}) \text{ as } |z| \to 0.$$

It is well known that a Padé approximation for e^{-z} exists, is unique and is representable by the formula $R_{p,q}(z) = P_{p,q}(z)/Q_{p,q}(z)$, where

$$P_{p,q}(z) = \Sigma_{j=0}^{p}\frac{(p+q-j)!p!(-z)^j}{(p+q)!j!(p-j)!},\ Q_{p,q}(z) = \Sigma_{j=0}^{q}\frac{(p+q-j)!q!z^j}{(p+q)!j!(q-j)!}.$$

In the papers [171]-[172] details of the location of poles and order of convergence of rational approximations in different regions are given.

Definition 3.3 *A rational approximation $r_{p,q}(\cdot) \in \pi_{p,q}$ for e^{-z} is said to be*
a) A-acceptable if $|r_{p,q}(z)| < 1$ for $Re(z) > 0$;
b) $A(\theta)$-acceptable if $|r_{p,q}(z)| < 1$ for $z \in \Sigma(\theta) = \{z : -\theta < arg(z) < \theta, z \neq 0\}$.

It is well known that $R_{q,q}(z), R_{q-1,q}(z)$ and $R_{q-2,q}(z)$ are A-acceptable. But for $q \ge 3$ and $p = q - 3$ the Padé functions are not A-acceptable.

Theorem 3.11 *[172] For any $q \ge 2$ and $p \ge 0$ the Padé approximation to e^{-z} has no poles in the sector $S_{p,q} = \{z : |arg(z)| < cos^{-1}\left(\frac{q-p-2}{p+q}\right)\}$, in particular for $p \le q \le p+4$ all poles lies in the left-half plane.*

Since $r(\cdot) \in \pi_{p,q}$ is an approximation of e^{-z} it is natural to construct an operator-function $r(\tau_n A_n)^k$ which could be considered as an approximation of $\exp(tA_n)$ for $t = k\tau_n$. We assume in this section for simplicity that $\|\exp(tA_n)\| \le M, t \in \overline{IR}_+$.

Theorem 3.12 *[41] Let condition (B) be satisfied. There is a constant C depending on r such that if r is A-acceptable, then*

$$\|r(\tau_n A_n)^k\| \le CM\sqrt{k} \text{ for } \tau_n > 0, k \in I\!N.$$

Remark 3.1 *The term $\sqrt{k}$ in Theorem 3.12 can not be removed in general, moreover, there are examples [52], [94] which shows that the inequality $\|r(\tau_n A_n)^k\| \ge c\sqrt{k}, k \in I\!N$, hold.*

We say that $r(\cdot) \in \pi_{p,q}$ is accurate of order $1 \le d \le p+q$ if $|e^{-z} - r(z)| = O(|z|^{d+1})$ as $|z| \to 0$.

Theorem 3.13 *[41] Let condition (B) be satisfied. Then there is a constant C depending on r such that, for r which is A-acceptable and accurate of order d, one gets*

$$\|r(\tau_n A_n)^k u_n^0 - \exp(tA_n)u_n^0\| \le CM\tau_n^d \|A_n^{d+1}u_n^0\| \text{ for } \tau_n > 0, k \in I\!N, u_n^0 \in D(A_n^{d+1}).$$

Theorem 3.14 *[41] Let condition (B_1) is satisfied. Then there is a constant C depending on r such that, for r which is A-acceptable and accurate of order d, one gets*

$$\|r(\tau_n A_n)^k u_n^0 - \exp(tA_n)u_n^0\| \le CM\tau_n^d \|A_n^{d}u_n^0\| \text{ for } \tau_n > 0, k \in I\!N, u_n^0 \in D(A_n^{d}).$$

Theorem 3.15 *[182], [159] Let condition (B_1) be satisfied. Then there is a constant C depending on r such that for r which is A-acceptable and accurate of order d with $|r(\infty)| < 1$ or condition (3.6) is satisfied, then one gets*

$$\|r(\tau_n A_n)^k u_n^0 - \exp(tA_n)u_n^0\| \le CM\frac{\tau_n^\gamma}{t^{d-\gamma}} \|A_n^{\gamma}u_n^0\| \text{ for } \tau_n > 0, 0 \le \gamma \le d, t = k\tau_n, k \in I\!N.$$

In [51] , [149], [151] the analogy of Theorems 3.13-3.15 was proved for multistep methods.

Let us recall that constant M_2 in condition (B_1) defines $0 < \alpha < \frac{\pi}{2}$ by $M_2 \sin\alpha < 1$ [107] such that

$$\|(\lambda I - A_n)^{-1}\| \le \frac{M}{|\lambda - \omega|} \text{ for any } \lambda \in \Sigma(\pi/2 + \alpha). \tag{3.8}$$

Theorem 3.16 *[52], [147] Let condition (B_1) be satisfied. Then there is a constant C depending on r such that, for r which is $A(\theta)$-acceptable, accurate of order d and $\theta \in (\pi/2 - \alpha, \pi/2]$ for α from condition (3.8), one gets*

$$\|r(\tau_n A_n)^k\| \le CM \text{ for } \tau_n > 0, k \in I\!N,$$

and

$$\|r(\tau_n A_n)^k - \exp(tA_n) - \gamma^k \exp(-\tau_n^{-b} a k_n (-A_n)^{-b})\| \le CM(k_n^{-d} + k_n^{-1/b}),\ t = k_n\tau_n,$$

where $\gamma = r(\infty)$ and a, b are some positive constants.

It is possible to show [148] that $\Pi_{j=1}^k r(\tau_{n,j}A_n)$ is a stable approximation for $\exp(\Sigma_{j=1}^k \tau_{n,j}A)$ with variable stepsize, but under condition $0 < c \le \tau_i/\tau_j \le C < \infty$, $i, j \in I\!N$.

3.3 Richardson's Extrapolation method

Let us consider the schemes (3.3) and (3.5) which have an order of convergence $O(\tau_n)$ and denote $\mathcal{U}_n^{\tau_n}(k_n) = U_n(t)u_n^0$ and $\overline{\mathcal{U}}_n^{\tau_n}(k_n) = \overline{U}_n(t)u_n^0, t = k_n\tau_n$. The following deferred approach to the limit is valid:

Theorem 3.17 *[164] Assume that condition (B) is satisfied. Then for* $\overline{V}_n(t) = 2\overline{\mathcal{U}}_n^{\tau_n}(k_n) - \overline{\mathcal{U}}_n^{\tau_n/2}(2k_n)$ *one has*

$$\|\overline{V}_n(t) - u_n(t)\| \le \tau_n^2 M e^{\omega t} t^2 \|A_n^3 u_n^0\|,\ t = k_n\tau_n.$$

If in addition the scheme (3.5) is stable, then for $V_n(t) = 2\mathcal{U}_n^{\tau_n}(k_n) - \mathcal{U}_n^{\tau_n/2}(2k_n), t = k_n\tau_n,$

$$\|V_n(t) - u_n(t)\| \le \tau_n^2 M e^{\omega t} t^2 \|A_n^3 u_n^0\|,\ t = k_n\tau_n.$$

Let us consider the Crank-Nicolson scheme

$$\frac{\tilde{U}_n(k\tau_n + \tau_n) - \tilde{U}_n(k\tau_n)}{\tau_n} = A_n \frac{\tilde{U}_n(k\tau_n + \tau_n) + \tilde{U}_n(k\tau_n)}{2},\ \tilde{U}_n(0) = I_n,\ k \in I\!N_0, \tag{3.9}$$

Theorem 3.18 *[164] Assume that condition (B) is satisfied and that scheme (3.9) is stable. Then* $\psi_n(t) = \frac{4}{3}\tilde{\mathcal{U}}_n^{\tau_n/2}(2k_n) - \frac{1}{3}\tilde{\mathcal{U}}_n^{\tau_n}(k_n)$ *satisfies*

$$\|\psi_n(t) - u_n(t)\| \le c\tau_n^4 e^{\omega t} t^2 \|A_n^6 u_n^0\|,\ t = k_n\tau_n.$$

In general, let us put $\mathcal{V}_n^{\tau_n}(t) = R_{p,q}(\tau_n A_n)^{k_n} u_n^0,\ t = k_n\tau_n.$

Theorem 3.19 *[164] Assume that condition (B) is satisfied,* $p = q$ *and the scheme which corresponds to* $\mathcal{V}_n^{\tau_n}$ *is stable. Then for* $\psi_n(t) = \frac{-1}{2^{2q}-1}\mathcal{V}_n^{\tau_n}(t) + \frac{2^{2q}}{2^{2q}-1}\mathcal{V}_n^{\tau_n/2}(t)$ *one gets*

$$\|\psi_n(t) - u_n(t)\| \le c\tau_n^{2q+2} e^{\omega t}\Big(\frac{t^{3/2}}{\sqrt{\tau_n}}\|A_n^{2q+3}u_n^0\| + t^3\tau_n^{2q-3}\|A_n^{4q+2}u_n^0\|\Big),\ t = k_n\tau_n.$$

Theorem 3.20 *[164] Assume that condition* (B_1) *is satisfied,* $p = q$ *and* $\tau_n\|A_n\| \le const.$ *Then for* $0 \le \gamma \le 2q$ *and* $\psi_n(t) = \frac{-1}{2^{2q}-1}\mathcal{V}_n^{\tau_n}(t) + \frac{2^{2q}}{2^{2q}-1}\mathcal{V}_n^{\tau_n/2}(t)$ *one gets*

$$\|\psi_n(t) - u_n(t)\| \le c\frac{\tau_n^{2q+2}}{t^{2q+2-\gamma}} e^{\omega t}\|A_n^\gamma u_n^0\|,\ t = k_n\tau_n.$$

3.4 Lax-type equivalence theorems with orders

The Lax equivalence Theorem on convergence of solution of approximation problem to the solution of the given well-posed Cauchy problem states that stability of the method is necessary and sufficient for convergence, provided it is consistent. More recently they furnish Lax's Theorem with orders, which make possible consider "not stable" approximations.

Definition 3.4 *The* C_0*-semigroups* $\exp(tA_n)$ *and* $\exp(tA)$ *are said to be consistent of order* $O(\varphi(\tau_n))$ *on a linear manifold* $\mathscr{U} \subset E$ *with respect to the semigroup* $\exp(\cdot A)$ *if* $\exp(tA)\mathscr{U} \subseteq D(A)$ *and there is a constant* C *such that*

$$\|(A_n p_n - p_n A)\exp(tA)x\| \le C\tau_n\varphi(\tau_n)e^{\omega t}|x|_{\mathscr{U}} \text{ for any } x \in \mathscr{U}, \tag{3.10}$$

where $|\cdot|$ *denotes the seminorm on* $\mathscr{U}$.

Definition 3.5 *The C_0-semigroups $\exp(tA_n)$ is said to be stable of order $O(M_n e^{\omega_n t})$ if there are constants M_n and ω_n such that*

$$\|\exp(tA_n)\| \le M_n e^{\omega_n t} \text{ for any } t \in \overline{I\!R}_+. \tag{3.11}$$

The following is a slight modification of [44]-[47] and [63]-[65], which was proved in [161].

Theorem 3.21 *Let the C_0-semigroup $\exp(\cdot A_n)$ be consistent of order $O(\varphi(\tau_n))$ on a linear manifold $\mathscr{U} \subset E$ with respect to the semigroup $\exp(\cdot A)$, $\exp(tA)\mathscr{U} \subset \mathscr{U}$ and $|\exp(tA)x|_{\mathscr{U}} \le M|x|_{\mathscr{U}}$. The following assertions are equivalent:*

(i) $\|\Big(\exp(tA_n)p_n - p_n\exp(tA)\Big)x\| \le 2M_n e^{\omega_n t} K(\frac{C_n}{2} t\varphi(\tau_n), x; E, \mathscr{U})$;

(ii) $\|\Big(\exp(tA_n)p_n - p_n\exp(tA)\Big)x\| \le M_n e^{\omega_n t}\begin{cases} M_x, & x \in E, \\ \frac{C_n}{2} t\varphi(\tau_n)|x|_{\mathscr{U}}, & t = k_n\tau_n \in [0,T], x \in \mathscr{U}; \end{cases}$

(iii) $\|\exp(tA_n)\| \le M_n e^{\omega_n t}$, $\|(A_n p_n - p_n A)\exp(tA)x\| \le C_n\tau_n\varphi(\tau_n)e^{\omega t}|x|_{\mathscr{U}}$ *for any* $x \in \mathscr{U}, t \in \overline{I\!R}_+$,

where M_x is a constant depending only on x and $K(t,x;E,\mathscr{U}) = \inf_{y\in\mathscr{U}}\Big\{\|x-y\|_E + t|y|_{\mathscr{U}}\Big\}$ is Peetre functional.

Definition 3.6 *The family of discrete semigroups $\{U_n(k_n\tau_n)\}$ is said to be consistent of order $O(\varphi(\tau_n))$ on a linear manifold $\mathscr{U} \subset E$ with respect to the semigroup $\exp(\cdot A)$ provided $\overline{\mathscr{U}} = E$ and*

$$\|(U_n(\tau_n)p_n - p_n\exp(\tau_n A))\exp(tA)x\| \le C\tau_n\varphi(\tau_n)|x|_{\mathscr{U}} \text{ for any } x \in \mathscr{U}. \tag{3.12}$$

Theorem 3.22 *Let $\exp(tA_n)\mathscr{U}_n \subset \mathscr{U}_n$, condition (B) holds and $|\exp(tA_n)x|_{\mathscr{U}_n} \le Ce^{\omega t}|x_n|_{\mathscr{U}_n}$ for any $x_n \in \mathscr{U}_n$ and $t > 0$. Then the following conditions are equivalent:*

(a) $\|(U_n(k_n\tau_n) - \exp(k_n\tau_n A_n))x_n\| \le M_n K(\frac{C_n k_n\tau_n}{2}\varphi(\tau_n), x_n, E_n, \mathscr{U}_n), n, k_n \in I\!N$;

(b) $\|(U_n(k_n\tau_n) - \exp(k_n\tau_n A_n))x_n\| \le M_n\begin{cases} M_{x_n}, & x_n \in E_n, \\ \frac{C_n}{2} t e^{t\omega}\varphi(\tau_n)|x_n|_{\mathscr{U}_n}, & x_n \in \mathscr{U}_n; \end{cases}$

(c) $\|U_n(k_n\tau_n)\|_{B(E_n)} \le M_n$, $\|(U_n(\tau_n) - \exp(\tau_n A))\exp(tA_n)x_n\| \le \frac{C_n M_n}{2}\tau_n e^{\omega t}\varphi(\tau_n)|x_n|_{\mathscr{U}_n}$, *where* $k_n\tau_n = t \in [0,T]$.

Definition 3.7 *The family of discrete semigroups $\{U(k_n\tau_n)\}$ is said to be stable of order $O(1/\psi(n^{-1}))$ if*

$$\|U_n(k_n\tau_n)\|_{B(E_n)} \le C/\psi(n^{-1}), \text{ for } n, k_n \in I\!N, 0 < \tau_n \le \tau^*, \tau_n k_n \in [0,T]. \tag{3.13}$$

Theorem 3.23 *Let the discrete semigroup $\{U(k_n\tau_n)\}$ be consistent of order $O(\varphi(\tau_n))$ on a linear manifold $\mathscr{U} \subset E$ with respect to the semigroup $\exp(\cdot A)$. The following assertions are equivalent:*

(i) $\|U_n(k_n\tau_n)\|_{B(E_n)} \le C/\psi(n^{-1})$;

(ii) $\|(U_n(k_n\tau_n)p_n - p_n\exp(k_n\tau_n A))x\| \le \frac{C}{\psi(n^{-1})}K(k_n\tau_n\varphi(\tau_n), x; E, \mathscr{U}), n, k_n \in I\!N$;

(iii) $\|(U_n(k_n\tau_n)p_n - p_n\exp(k_n\tau_n A))x\| \le \frac{C}{\psi(n^{-1})}\begin{cases} M_x, & x \in E, \\ k_n\tau_n\varphi(\tau_n)|x|_{\mathscr{U}}, & k_n\tau_n \in [0,T], x \in \mathscr{U}, \end{cases}$

where M_x is a constant depending only on x.

Theorem 3.24 *Let $|\exp(tA)x|_{\mathscr{U}} \le C|x|_{\mathscr{U}}$ for any $x \in \mathscr{U}$ and $t \in [0,T]$. Then the following conditions are equivalent:*

(i) The family of operators $\{U(k_n\tau_n)\}$ is consistent of order $O(\varphi(\tau_n))$ on a linear manifold $\mathscr{U} \subset E$ with respect to the semigroup $\exp(\cdot A)$ and stable of order $O(1/\phi(n^{-1}))$;

(ii) $\|(U_n(k_n\tau_n)p_n - p_n\exp(k_n\tau_n A))x\| \le \frac{C}{\psi(n^{-1})}K(k_n\tau_n\varphi(\tau_n), x; E, \mathscr{U}), n, k_n \in I\!N$;

(iii) $\|(U_n(k_n\tau_n)p_n - p_n\exp(tA))x\| \le \frac{C}{\psi(n^{-1})}\begin{cases} M_x, & x \in E, \\ k_n\tau_n\varphi(\tau_n)|x|_{\mathscr{U}}, & t = k_n\tau_n \in [0,T], x \in \mathscr{U}. \end{cases}$

On the extension of Lax-Richtmyer theory see [154], [178].

For the particular case, when $E = L^p(I\!R^d)$ and operator $A \equiv P(D) = \Sigma_{|\alpha|\le r} p_\alpha D^\alpha$ in E, one the can consider Cauchy problem

$$\frac{\partial u(x,t)}{\partial t} = P(D)u(x,t),\ u(x,0) = u^0(x), x \in \overline{I\!R}_+, \tag{3.14}$$

with $P(D)$ such that (3.14) is well-posed in the sense $\|u(\cdot,t)\|_{L^p(I\!R)} \le c\,\|u^0(\cdot)\|_{L^p(I\!R)}, t \in \overline{I\!R}_+$.

Let us denote $\hat{P}(\xi) = \Sigma_{|\alpha|\le r} p_\alpha (i\xi)^\alpha$. It is well-known that (3.14) well-posed iff $\|\exp(t\hat{P})\|_{M_p} \le C$, $t \in \overline{I\!R}_+$, where M_p is a space of Fourier multipliers.

Semidiscrete approximation of (3.14) is given by

$$\frac{\partial u_n(x,t)}{\partial t} = P_h(D)u_n(x,t),\ u_n(x,0) = u_n^0(x), x \in \overline{I\!R}_+, \tag{3.15}$$

where $P_h(D_h) = h^{-r}\Sigma_{|\alpha|\le r} p_\alpha(h)\Sigma_{\beta\in I_\alpha} b_\beta u_n(x+\beta h,t)$ and $\hat{P}_h(\xi) = h^{-r}\Sigma_{|\alpha|\le r} p_\alpha(h)\Sigma_{\beta\in I_\alpha} b_\beta e^{i\langle\xi,h\beta\rangle}$. Operator $P_h(D_h)$ is said to be consistent with operator $P(D)$ with order μ if $\hat{P}_h(\xi) - \hat{P}(\xi) = h^\mu|\xi|^{r+\mu}Q(h\xi), r = deg\hat{P}_h(\xi), Q$ is an infinitely differentiable function and $|Q(\eta)| \ge Q_0 > 0$ for $0 < |\eta| \le \epsilon_0$.

Theorem 3.25 *[40] Let $P(D)$ and $P_h(D_h)$ be consistent of order μ and (3.14) and (3.15) be well-posed. Then for every $T > 0$ there exists $C > 0$ such that $\|\left(e^{tP_h(D_h)} - \exp(tP(D))\right)u^0\|_{L^2(I\!R^d)} \le ch^\mu\|u^0\|_{W^{2,r+\mu}(I\!R^d)}$ and for $0 < s < r + \mu$*

$$\begin{aligned}
\|\left(e^{tP_h(D_h)} - \exp(tP(D))\right)u^0\|_{L^2(I\!R^d)} &\le ch^{\frac{s\mu}{\mu+r}}\|u^0\|_{B_2^s},\\
\|\left(e^{tP_h(D_h)} - \exp(tP(D))\right)u_n^0\|_{L^\infty(I\!R^d)} &\le ch^{\frac{s\mu}{\mu+r}}\|u^0\|_{B_2^{d/2+s}},
\end{aligned}$$

where $B_p^\theta = B_{p,\infty}^\theta$ is a Besov space.

It is remarked in [29], that, for the quite general case $B_{p,q}^\theta = (L^p(I\!R), D(A))_{\theta,q}$.

If we consider full discretization scheme for (3.14) in the form $L_h U_n^{k+1} = B_h U_n^k, k = 0,1,2,\ldots$, where $L_h v = \Sigma_\beta a_\beta(h)v(x+\beta h), B_h v = \Sigma_\beta b_\beta(h)v(x+\beta h)$, then a discrete semigroup can be constructed as $U_n(k\tau_n)u_n^0 = \mathscr{F}^{-1}\left(\hat{U}_n^k(\xi)\hat{u}_n^0\right), \hat{U}_n(\xi) = \hat{B}_n(\xi)/\hat{L}_n(\xi), \hat{B}_n(\xi) = \Sigma_\beta a_\beta(h)e^{\langle\xi,\beta h\rangle}$ (the time step τ_n is tied to h by $\tau_n/h^r = constant$). Such a finite difference operator, $U_n(k\tau_n)$, approximates (3.14) with order μ if $\hat{U}_n(\xi) = e^{\tau_n\hat{P}(\xi)} + O(h^{r+\mu} + |\xi|^{r+\mu})$ as $\xi, h \to 0$.

Theorem 3.26 *[40] Let (3.14) be well-posed and $U_n(k\tau_n)$ be stable in $E = L^2(I\!R^d)$ and approximate (3.14) with order $\mu > 0$. Then for any $T > 0$ there is a constant $c > 0$ such that*

$$\|\left(U_n(t) - \exp(tP(D))\right)u_n^0\|_{L^2(I\!R^d)} \le ch^\mu\|u_n^0\|_{W^{2,r+\mu}(I\!R^d)}$$

and for $0 < s < r + \mu$

$$\begin{aligned}
\|\left(U_n(t) - \exp(tP(D))\right)u^0\|_{L^2(I\!R^d)} &\le ch^{\frac{s\mu}{\mu+r}}\|u^0\|_{B_2^s},\\
\|\left(U_n(t) - \exp(tP(D))\right)u^0\|_{L^\infty(I\!R^d)} &\le ch^\mu\|u^0\|_{B_{2,1}^{d/2+\mu+r}},\\
\|\left(U_n(t) - \exp(tP(D))\right)u^0\|_{L^\infty(I\!R^d)} &\le ch^{\frac{s\mu}{\mu+r}}\|u^0\|_{B_2^{d/2+s}},\ t = k\tau_n \in [0,T].
\end{aligned}$$

Inverse results: order of convergence implies smoothness of u_n^0, see [29], [40].

Time discretization of parabolic problems with memory by backward Euler method was considered in [24]. Stability and error estimates take place in a Banach space framework, and the results are used to derive error estimates in the L_2 and maximum norms for piecewise linear finite-element discretizations in two space dimensions.

4 Backward Cauchy problem

Let us consider in the Banach space E the backward Cauchy problem:

$$\begin{aligned} &v'(t) = Av(t),\ t \in [0,T], \\ &v(T) = v^T, \end{aligned} \tag{4.1}$$

where we are going to find element $v(0)$. At least in two important cases it is not a well-posed problem, namely, if A is unbounded and generates an analytic C_0-semigroup or if the C_0-semigroup $\exp(\cdot A)$ is compact. Indeed, in these situations the problem $\exp(TA)x = v^T$ is ill-posed [48], [97], [193] in the sense that the operator $\exp(-TA)$ is not bounded on E and, moreover, in general $D(\exp(-TA)) \neq E$. This means that the Cauchy problem (4.1) has a solution in general only for some (but not every) initial data v^T and the solution $v(0)$, if it is exists, does not depend continuously on the initial data. After changes of variables and putting $v(\eta) = u(T-\eta)$ one can rewrite the problem (4.1) in the form

$$\begin{aligned} &u'(t) = -Au(t),\ t \in [0,T], \\ &u(0) = u^0, \end{aligned} \tag{4.2}$$

where $u^0 = v^T$ is given and $u(T)$ is the element which we are going to find. In this section we are going consider the approximation of (4.2) with operator A, which generates analytic C_0-semigroup.

Definition 4.1 *A bounded linear operator $R_{\epsilon,T}$ on the space E is called a regularizator for the Cauchy problem (4.2) if for any $\delta > 0$ and any $u^0 \in E$, for which a solution of (4.2) exists, there exists $\epsilon = \epsilon(\delta) > 0$ such that $\epsilon(\delta) \to 0$ as $\delta \to 0$ and $\sup_{\|u^\delta - u^0\| \leq \delta} \|R_{\epsilon(\delta),T} u^\delta - \exp(-TA)u^0\| \to 0$ as $\delta \to 0$.*

In [137] it is proved that for the existence of a linear regularizator of the problem (4.2), which commute with operator A, it is necessary and sufficient that $-A$ generates C_ϵ-semigroups $\mathcal{S}_\epsilon(t)$, $0 \leq t \leq T$, such that C_ϵ strongly converges to the identity operator I as $\epsilon \to 0$.

There are a lot of regularizators, which could be considered for problem (4.2). For example, in [162] it was shown that if $-A_n^2$ generates a cosine operator function, then the method of quasireversibility, which is given by the Cauchy problems

$$u'_{n,\alpha}(t) = -A_n u_{n,\alpha}(t) - \alpha A_n^2 u_{n,\alpha}(t),\ u_{n,\alpha}(0) = u_n^0,$$

is a regularization method for (4.2), and $\|u_{n,\alpha}(T) - p_n u(T)\| \leq C\alpha\Big(\|u_n^0 - p_n u^0\|/\delta + \rho\Big)$, where $\alpha = \alpha(\delta) = 1/\Big(\ln(1/\delta) - \ln\ln(1/\delta) - o(\ln^{-1}(1/\delta))\Big)$. In this case $\mathcal{S}_\alpha(t) \equiv \exp(-tA)\exp(-\alpha T A^2)$ is a C_α-semigroup with $C_\alpha = \exp(-\alpha T A^2)$ and $C_\alpha \to I$ as $\alpha \to 0$. Moreover, the generator of this C_α-semigroup is $-A$.

It has been shown in [57] that the stochastic differential equation

$$\begin{aligned} &du_\alpha(t) = -Au_\alpha(t)dt - \alpha A u_\alpha(t) dw(t), \\ &u(0) = u^0, \end{aligned} \tag{4.3}$$

where $w(\cdot)$ is a standard one-dimensional Wiener process, yields a stochastic regularization of (4.2). Explicitly, the operator-function

$$U_\alpha(t)u^0 = \frac{1}{2\pi i}\int_\Gamma e^{-t\lambda - \alpha\Big(w(t)-w(0)\Big)\lambda - \frac{1}{2}\alpha^2\lambda^2|t|}(\lambda - A)^{-1}u^0 d\lambda,\ t > 0,$$

which represents a solution of (4.3) for any $u^0 \in \mathfrak{A}_c(A)$, possesses the following properties:

$$\lim_{\alpha \to 0} \|U_\alpha(T)u^0 - \exp(-TA)u^0\| = 0, \tag{4.4}$$

$$\|U_\alpha(t)\| \le \frac{c_1}{\alpha\sqrt{|t|}} \exp\left(c_2\frac{\sqrt{|t|}}{\alpha} + c_3|t|^{-\mu})^2\right) + b(\alpha,|t|) \text{ for any } \alpha > 0. \tag{4.5}$$

Here the function $b(\alpha,t)$ is bounded in the parameters α and t and $\mathfrak{A}_c(A)$ is the set of entire vectors of operator A. This means that by virtue of the inequality

$$\|U_\alpha(T)u^\delta - \exp(-TA)u^0\| \le \|U_\alpha(T)\|\,\|u^\delta - u^0\| + \|U_\alpha(T)u^0 - \exp(-TA)u^0\|, \tag{4.6}$$

there is a dependence on $\alpha = \alpha(\delta)$ in such a way that $U_\alpha(T)$ becomes a regularizator. The operator-function $t \mapsto \exp\big((T-t)A\big)U_\alpha(T)$, $0 \le t \le T$, is a C_α-semigroup with $C_\alpha = \exp(TA)U_\alpha(T)$. One can see that $C_\alpha \to I$ as $\alpha \to 0$ and that the generator of this C_α-semigroup is $-A$.

4.1 C-semigroups and ill-posed problems

Let C be a bounded linear operator on the Banach space E, i.e. $C \in B(E)$, and let $T > 0$ be some finite number.

Definition 4.2 *[188]. A family of bounded operators $\{\mathcal{S}(t) : 0 \le t < T\}$ is called a local C-semigroup on E if*
(i) $\mathcal{S}(t+s)C = \mathcal{S}(t)\mathcal{S}(s)$ for $t, s, t+s \in [0,T)$,
(ii) $\mathcal{S}(0) = C$,
(iii) $\mathcal{S}(\cdot)$ is strongly continuous on $[0,T)$.

Clearly, $\mathcal{S}(\cdot)$ is a commutative family. A local C-semigroup is called nondegenerate if condition $\mathcal{S}(t)x = 0$ for all $t \in (0,T)$ implies $x = 0$. It is seen from Definition 4.2 that a local C-semigroup is nondegenerate [60] if and only if C is injective, i.e., $\mathcal{N}(C) = \{0\}$. Concerning construction with $\mathcal{N}(C) \ne \{0\}$ see [109], [110]. It is very interesting question how to apply the case of noninjective C, i.e. degenerate C-semigroups, for ill-posed problems. Unfortunately this approach still is not realized.

Starting from now on we will consider only the case that $C \in B(E)$ is an injective operator.

Definition 4.3 *The generator of $\{\mathcal{S}(t) : 0 \le t < T\}$ is defined as the limit $-Gx := C^{-1}\lim_{h\to 0+}\frac{1}{h}(\mathcal{S}(h)x - Cx), x \in D(G)$, with a natural domain $D(G) := \Big\{x \in E : \exists \lim_{h\to 0+}\frac{1}{h}(\mathcal{S}(h)x - Cx) \in R(C)\Big\}$.*

Proposition 4.1 *[179] The operator G is closed, $R(C) \subseteq \overline{D(G)}$ and $C^{-1}GC = G$.*

We denote the C-semigroup $\mathcal{S}(\cdot)$ with generator $-G$ by $\mathcal{S}(\cdot G)$. Next, let $\tau \in (0,T)$. Put

$$L_\tau(\lambda)x := \int_0^\tau e^{-\lambda t}\mathcal{S}(tG)x\,dt,\ x \in E, \lambda > 0. \tag{4.7}$$

This is the so-called *local Laplace transform* of $\mathcal{S}(\cdot G)$.

Proposition 4.2 *Let $\mathcal{S}(\cdot G)$ be a local C-semigroup and $L_\tau(\cdot)$ be the local Laplace transform of $\mathcal{S}(\cdot G)$. Then for any $x \in E$ one has $L_\tau(\lambda)x \in D(G)$ and*

$$(\lambda + G)L_\tau(\lambda)x = Cx - e^{-\lambda\tau}\mathcal{S}(\tau G)x \text{ for all } \tau \in [0,T) \text{ and } \lambda > 0. \tag{4.8}$$

In the case of local C-semigroups spectrum $\sigma(-G)$ could be located on the line $[0,\infty)$. So in this case in general the Laplace transform of the local C-semigroup does not exist in principle and we follow the ideas from [27], [179], [188]. The function $L_\tau(\lambda)$ with the property (4.8) is said to be asymptotic resolvent.

Theorem 4.1 *[179] Let A be a closed linear operator in E and let $C \in B(E)$ be injective.*

(i) If operator A is the generator of a local C-semigroup $\{S(t); 0 \leq t < T\}$ on E, then there exists an asymptotic C-resolvent $L_\tau(\lambda)$ of $-A$ such that

$$\|\frac{d^m}{d\lambda^m} L_\tau(\lambda)x\| \leq M_\tau \frac{m!}{\lambda^{m+1}}\|x\|,\ x \in E, \tag{4.9}$$

with $0 \leq m/\lambda \leq \tau, \lambda > a, m \in I\!N \cup \{0\}$, and operator A satisfies $C^{-1}AC = A$.

(ii) If $-A$ has an asymptotic resolvent which satisfies (4.9), and $CD(A)$ is dense in $D(A)$, $D(C^{-1}AC) \subset D(A)$, i.e. $Cx \in D(A)$ and $ACx \in R(C)$ imply $x \in D(A)$, then the part A_0 of A in $E_0 := \overline{D(A)}$ generates a local C-semigroup on E_0, with C equal to $C_0 := C|_{E_0}$.

In particular, under the assumption that $\overline{CD(A)} = E$, the operator $-A$ generates a local C-semigroup on E if and only if $C^{-1}AC = A$ and there exists an asymptotic C-resolvent satisfying (4.9). In this case, A has dense domain.

Remark 4.1 *An asymptotic C-resolvent $L_\tau(\lambda)$ of operator $-A$ is compact for some $\lambda \in \mathbb{C}$ (and then for any λ large enough) if and only if $\mathcal{S}(\cdot A)$ is compact and uniformly continuous in t. Indeed, if $\mathcal{S}(\cdot A)$ is compact then by (4.7) and [217] it follows that $L_\tau(\lambda)$ is compact. Conversely, taking derivative of $L_\tau(\lambda)$ in τ and using the fact that $\mathcal{S}(\cdot A)$ is uniformly continuous in t we have that $\mathcal{S}(\cdot A)$ is compact as the uniform limit of compact operators. This fact could be used in the approximation of semilinear equations in case of the C-semigroups approach (cf. section 6).*

Let us consider the abstract Cauchy problem, which is given by (4.2).

Definition 4.4 *A function $u(\cdot)$ is called a solution to (ACP; T,y) if $u(\cdot)$ is continuously differentiable in $t \in [0,T)$, $u(t) \in D(A)$ for all $0 \leq t < T$, and $u(\cdot)$ satisfies (4.2). We denote (ACP;T,y) with $y \in CD(A)$, also by (ACP;T,CD(A)).*

Definition 4.5 The Cauchy problem $(ACP;T,CD(A))$ is said to be *generalized well-posed* if for every $y \in CD(A)$, there is a unique solution $u(\cdot;y)$ to $(ACP;T,y)$ such that $\|u(t;y)\| \leq M(t)\|C^{-1}y\|$ for $0 \leq t < T$ and $y \in CD(A)$, where the function $M(t)$ is bounded on every compact subinterval of $[0,T)$.

We have to stress here that generalized well-posedness in the sense of Definition 4.5 is more general than in the case of the problem in (3.1). Moreover, we can state that this generalized well-posedness is a solvability condition of (4.2) for which a regularizator exists.

Theorem 4.2 *[179] Let C be a bounded linear injection on E and let A be a closed linear operator. Then the following assertions are equivalent:*

(I) The operator $-A$ is the generator of a local C-semigroup;

(II) $C^{-1}AC = A$, and the problem: $v'(t) = -Av(t) + Cx$, $t \in [0,T), v(0) = 0$, has a unique solution for every $x \in X$.

If either $\rho(A) \neq \emptyset$ or A has dense domain, (I) and (II) are also equivalent to

(III) $C^{-1}AC = A$, and the problem $\left(ACP;T,CD(A)\right)$ is generalized well-posed. Moreover, $u(t;y) = C^{-1}\mathcal{S}(tA)y$, $t \in [0,T)$, is a unique solution for every initial value $y \in CD(A)$.

Since the local C-semigroups are regularizators of the ill-posed problem (4.2) it is very important to present approximation theory of local C-semigroups.

4.2 Semidiscrete approximation theorem

Let us consider, within the general discretization scheme, the semidiscrete approximation of the problem (4.2) in the Banach spaces E_n :

$$\begin{aligned} &u_n'(t) = -A_n u_n(t), \quad t \in [0,T), \\ &u_n(0) = u_n^0, \end{aligned} \tag{4.10}$$

where the operators $-A_n$ are generators of local C_n-semigroups, which are compatible with the operator $-A$ and $u_n^0 \to u^0$. We understand compatibility in the sense of the general approximation scheme as the $\mathcal{PP}$-convergence of $C_n \to C$ and the $\mathcal{PP}$-convergence of resolvents $(\tilde{\lambda} - A_n)^{-1} \to (\tilde{\lambda} - A)^{-1}$ for some $\tilde{\lambda} \in \rho(A) \cap \rho(A_n)$. Recall that in our general case (4.1) such $\tilde{\lambda}$ exists, since the conditions (A) and (B) from Theorem 3.1 are naturally assumed to be satisfied.

Theorem 4.3 *[211] (*Theorem ABC-C*) Under the assumption $\overline{CD(A^2)} = E$, the following conditions (A_c) together with (B_c) are equivalent to condition (C_c).*

(A_c) Consistency. $C_n \to C$ and operators A_n, A are compatible;

(B_c) Stability. For any $0 < \tau < T$ there is some constant M_τ, not depending on n, such that

$$\|\mathcal{S}(tA_n)\| \leq M_\tau \text{ for } 0 \leq t \leq \tau \text{ and } n \in I\!N;$$

(C_c) Convergence. For any $0 < \tau < T$ we have $\max_{t\in[0,\tau]} \|\mathcal{S}(tA_n)x_n^0 - p_n\mathcal{S}(tA)x^0\| = 0$, as $n \to \infty$, whenever $x_n^0 \to x^0$.

Remark 4.2 *In the case of exponentially bounded C-semigroups [61], [62] we can trivially change the condition (A_c) to condition*

(A') $C_n \to C$ and $(\tilde{\lambda} - A_n)^{-1}C_n \to (\tilde{\lambda} - A)^{-1}C$ for some $\tilde{\lambda} \in \mathbb{C}$; see [222] for details. Since, the construction can be done just as with condition (A'), in this case we do not need to assume that $(\tilde{\lambda} - A_n)^{-1} \to (\tilde{\lambda} - A)^{-1}$ for some $\tilde{\lambda}$.

Remark 4.3 *We have put the condition $\overline{CD(A^2)} = E$ for simplicity. For the general case one gets convergence on the set $\overline{CD(A^2)}$. In the case of integrated semigroups such situations have been well investigated, see for instance [30], [32]. Actually the paper [30] is devoted to the following effect observed in the study of convergence of semigroups. Suppose we are given a sequence of uniformly bounded semigroups $\{\exp(tA_n), t \geq 0\}, n \geq 1$ ($\|\exp(tA_n)\| \leq M, t \in \overline{I\!R}_+$) acting in the Banach space E. Assume furthermore that the limit $\lim_{n\to\infty}(\lambda I - A_n)^{-1}x = S(\lambda)x$ exists for any $x \in E$. If*

$$\overline{\mathcal{R}(S(\lambda))} = E \tag{4.11}$$

($\mathcal{R}(S(\lambda))$ is common for all $\lambda > 0$), the semigroups in question converge strongly, by the Trotter - Kato theorem. One may also show that if condition (4.11) is relaxed, the limit

$$\lim_{n\to\infty} \exp(tA_n)x \tag{4.12}$$

exists for all $x \in \overline{\mathcal{R}(S(\lambda))}$ (see e.g. [112], [67] p. 34, or [30], [32]). As observed by T.G. Kurtz [112] for any $x \in E$ there exists the limit

$$\lim_{n\to\infty} \int_0^t \exp(sA_n)x\,ds. \tag{4.13}$$

In general, however, one cannot expect that (4.12) holds for $x \notin \overline{\mathcal{R}(S(\lambda))}$. This effect is of course related to Arendt's theorem, or rather to generation theorem for "absolutely continuous integrated semigroups" presented in [34].

Let us consider a semi-discretization of problem (4.3) in Banach spaces E_n

$$\begin{aligned} du_{n,\alpha}(t) &= -A_n u_{n,\alpha}(t)dt - \alpha A_n u_{n,\alpha}(t)dw(t), \\ u_{n,\alpha}(0) &= u_n^0, \end{aligned} \tag{4.14}$$

where $u_n^0 \to u^0$, the operators A_n generate analytic semigroups and $\{(\Omega, \mathcal{F}, I\!P), w(t)\}$ is the standard one-dimensional Wiener process (Brownian motion). As usual the symbol $I\!E[\cdot]$ will denote mathematical expectation.

We emphasize that we consider the situation where $\sigma(A_n)$, $\sigma(A) \subset \mathbb{C} \setminus \Sigma(\frac{3}{4}\pi)$.

Theorem 4.4 *[211] Let the conditions (A) and (B_1) of Theorem 3.2 be satisfied and let $\delta_n > 0$ be a sequence which converges to 0 if $n \to \infty$. Then there exists a sequence α_n such that $u_{n,\alpha_n}(t) \to u(t)$ for every $t \in [0,T]$ as $n \to \infty$. Here $u_{n,\alpha_n}(\cdot)$ is a solution of (4.14) and $u(\cdot)$ is a solution of (4.2) with $u^0 \in \mathfrak{A}_c(A)$. Convergence is to be understood in the following sense:*

$$\sup_{\|u_n^0 - p_n u^0\| \le \delta_n} \|u_{n,\alpha_n}(t) - p_n u(t)\| \to 0, \quad I\!P\text{-almost surely as } \delta_n \to 0.$$

4.3 Approximation by discrete C-semigroups

Following Section 3 we denote by $\{T_n(\cdot)\}$ a family of discrete semigroups, on E_n, respectively, i.e., $T_n(t) = T_n(\tau_n)^{k_n}$, where $k_n = [t/\tau_n]$. We define the generator of $T_n(\cdot)$ by the formula $-A_n = \dfrac{1}{\tau_n}(T_n(\tau_n) - I)$ and we are interested in the process $\tau_n \to 0, k_n, n \to \infty$. We assume that $C_n \in B(E_n)$ is an injective operator such that $T_n C_n = C_n T_n$. The discrete C_n-semigroup $U_n(\cdot)$ is defined as $U_n(t) = T_n(t)C_n$. In this section we also assume that the dimension of each space E_n is finite, but $\dim(E_n) \to \infty$ as $n \to \infty$.

Theorem 4.5 *[211] (Theorem ABC-C-discr) Under condition (A) in Theorem 3.1 and the assumption $\overline{CD(A^2)} = E$, the following conditions (A_{cd}) and (B_{cd}) together are equivalent to condition (C_{cd}).*

(A_{cd}) Consistency. $C_n \to C$, operators A_n, A are compatible and $A_n \in B(E_n)$, $n \in I\!N$;

(B_{cd}) Stability. For any $0 < \tau < T$ there is some constant M_τ, not depending on n, such that

$$\|U_n(t)\| \le M_\tau \text{ for all } 0 \le t \le \tau < T \text{ and } n \in I\!N.$$

is satisfied uniformly for any choice of $\{\tau_n\}$ and $\{k_n\}$ as long as $\tau_n \to 0$, and $k_n = [t/\tau_n]$;

(C_{cd}) Convergence. For any $0 < \tau < T$ one gets $\max_{t\in[0,\tau]} \|U_n(t)x_n^0 - p_n \mathcal{S}(tA)x^0\| \to 0$ as $\tau_n \to 0, n \to \infty$, whenever $x_n^0 \to x^0$.

Theorem 4.6 *[211] Let the conditions (A_c) and (B_c) be satisfied. Assume that condition (A) in Theorem 3.1 and the assumption $\overline{CD(A^2)} = E$ are satisfied and also that $\tau_n \|A_n^2 C_n^{-1}\| \le \frac{q}{M_\tau T}$ with $q < 1$. Then*

$$\|U_n(t)\| \le M_\tau (1-q)^{-1} \text{ for } 0 \le t \le \tau < T \text{ and any } n \in I\!N.$$

is satisfied uniformly for any choice of $\{\tau_n\}$ and $\{k_n\}$ with $\tau_n \to 0$, as long as $k_n = \left[\frac{t}{\tau_n}\right]$. Moreover, for any $0 < \tau < T$ one gets $\max_{t\in[0,\tau]} \|U_n(t)x_n^0 - p_n \mathcal{S}(tA)x^0\| \to 0$ as $\tau_n \to 0$, $n \to \infty$, whenever $x_n^0 \to x^0$.

Remark 4.4 *As a matter of fact the scheme $\overline{U}_n(t) \equiv (I + \tau_n A_n)^{-k_n} C_n$ with $t = k_n \tau_n$ could be constructed even under condition (3.4). Indeed, $\tau_n A_n = \tau_n \lambda A_n (\lambda - A_n)^{-1} - \tau_n A_n^2 (\lambda - A_n)^{-1}$, and by the choice of λ we can make the second term less than ϵ and then by choosing τ_n appropriately for a fixed λ, we obtain $\|\tau_n A_n\| \le 2\epsilon$, so that the scheme $\overline{U}_n(\cdot)$ is well defined.*

Remark 4.5 *In contrast to the well-posed case, for ill-posed problems it looks that the implicit and explicit methods of discretization in time are not so different in the sense of stability advantages (compare with Theorems 3.5, 3.6). Moreover, under condition (3.4) from the identity*

$$(I-\tau_n A_n)^{k_n} C_n = (I-\tau_n^2 A_n^2)^{k_n}(I+\tau_n A_n)^{-k_n} C_n$$

and inequality $\|(I \pm \tau_n^2 A_n^2)^{k_n}\| \le C e^{t\tau_n \|A_n^2\|}$, $t = k_n\tau_n$, *it follows that the stability properties of the implicit and explicit methods are the same.*

There are a lot of stochastic finite-difference schemes which could be written for problem (4.14). For example, some of the simplest are

$$U_{n,\alpha}(t+\tau_n) - U_{n,\alpha}(t) = -\tau_n A_n U_{n,\alpha}(t) - \alpha\,\Delta w(t) A_n U_{n,\alpha}(t), \tag{4.15}$$

$$\bar{U}_{n,\alpha}(t+\tau_n) - \bar{U}_{n,\alpha}(t) = -\tau_n A_n \bar{U}_{n,\alpha}(t+\tau_n) - \alpha\,\Delta w(t) A_n \bar{U}_{n,\alpha}(t), \tag{4.16}$$

where $\Delta w(t) = \Big(w(t) - w(t-\tau_n)\Big)$, $t = k_n\tau_n$, and $U_{n,\alpha}(0) = \bar{U}_{n,\alpha}(0) = I_n$.

Theorem 4.7 *[211] Let the conditions (A) and (B_1) of Theorem 3.2 be satisfied. Assume that the stability conditions (3.4) and* $\tau_n \|A_n^2\| e^{c\|A_n\|} = O(1)$ *are fulfilled for some constant* $c > 0$. *Then for* $\alpha_n = \sqrt{\tau_n}$ *the scheme (4.15) behaves stably in the following sense:*

$$\eta_n := \sup_{n\in\mathbb{N}} \sup \left\{ I\!E \left[\left\| U_{n,\alpha_n}(t) u_n^0 - \exp\left(-tA_n + \alpha_n (w(t) - w(0)) A_n - \frac{t}{2}\alpha_n^2 A_n^2 \right) u_n^0 \right\| \right] : \|u_n^0\| \le 1 \right\} \to 0,$$

and converges in the following sense:

$$I\!E\left[\|U_{n,\alpha_n}(t) u_n^0 - p_n u(t)\| \right] \le C\sqrt{\tau_n} \|A \exp(-TA) u^0\| + \|u_{n,\alpha_n}(t) - p_n u_{\alpha_n}(t)\| + C\eta_n \|u_n^0\|, \quad 0 < t \le T.$$

For the scheme $\overline{U}_{n,\alpha_n}(\cdot)$ similar notions are employed.

We can also study the convergence behaviour of more sophisticated numerical methods. For example, in [42] in order to approximate (4.14) the following Runge-Kutta scheme was considered

$$\begin{aligned} Y_1 &= U_{n,\alpha}(t) + \sqrt{\tau_n}\alpha A_n U_{n,\alpha}(t), \\ U_{n,\alpha}(t+\tau_n) - U_{n,\alpha}(t) &= -\tau_n A_n U_{n,\alpha}(t) + \alpha \Delta w(t) A_n U_{n,\alpha}(t) + \\ &+ \frac{\sqrt{\tau_n}}{2}\left(\frac{(\Delta w(t))^2}{\sqrt{\tau_n}} - 1 \right)\Big(\alpha_n A_n Y_1 - \alpha A_n U_{n,\alpha}(t) \Big). \end{aligned} \tag{4.17}$$

Thus the solution can be written in the form

$$\begin{aligned} U_{n,\alpha}(t+\tau_n) = \\ = (I_n - \tau_n A_n - \frac{\alpha^2\tau_n}{2} A_n^2)^{k_n} \Pi_{k=1}^{k_n} \Big(I_n + Z_n^{-1}\alpha \Delta w(t) A_n + (Z_n^{-1}/2)\alpha^2 \Delta w(t)^2 A_n^2 \Big) U_{n,\alpha}(0), \end{aligned} \tag{4.18}$$

where $Z_n = (I_n - \tau_n A_n - \frac{\alpha^2\tau_n}{2} A_n^2)$.

Theorem 4.8 *[42] Let the conditions (A) and (B) of Theorem 3.2 be satisfied. Assume that the stability conditions (3.4) and* $\tau_n \|A_n^2\| e^{c\|A_n\|} = O(1)$ *are fulfilled for some constant* $c > 0$. *Then for* $\alpha_n = \sqrt{\tau_n}$ *the scheme (4.18) behaves stably in the following sense:*

$$\eta_n := \sup \left\{ I\!E \left[\left\| U_{n,\alpha_n}(t) u_n^0 - e^{\left(-tA_n + \alpha_n (w(t) - w(0)) A_n - \frac{t}{2}\alpha_n^2 A_n^2 \right)} u_n^0 \right\| \right] : \|u_n^0\| \le 1 \right\} \to 0,$$

and converges in the following sense:

$$I\!E\left[\|U_{n,\alpha_n}(t) u_n^0 - p_n u(t)\| \right] \le$$

$$\le C\sqrt{\tau_n} \|A \exp(-TA) u^0\| + \|u_{n,\alpha_n}(t) - p_n u_{\alpha_n}(t)\| + C\eta_n \|u_n^0\|, \quad 0 < t \le T.$$

In the case of the well-posed problem

$$\begin{cases} du_\alpha(t) = Au_\alpha(t)dt + \alpha Au_\alpha(t)dw(t), \\ u(0) = u^0, \end{cases} \tag{4.19}$$

where the operator A generates an analytic C_0-semigroup, the semidiscrete and full-discretisation schemes do not need additional stability assumptions and the order of convergence will be defined just by the consistency property of the scheme. More exactly the term $e^{\lambda t}$ under the integral leads to the absolute convergence of the integral independently of the behaviour of α on any compact set. For example we have the following

Theorem 4.9 *[42] Let the conditions (A) and (B") of Theorem 3.2 be satisfied. Assume that the stability conditions (3.4) is fulfilled for some constant $C > 0$. Then for any $\alpha_n \in [0, \alpha']$ the scheme like (4.18) behaves stably in the following sense:*

$$\sup \left\{ I\!E \left[\left\| U_{n,\alpha_n}(t)u_n^0 - e^{\left(tA_n + \alpha_n(w(t)-w(0))A_n - \frac{t}{2}\alpha_n^2 A_n^2\right)} u_n^0 \right\| \right] : \|u_n^0\| \le 1 \right\} \le \tau_n,$$

and converges in the following sense:

$$I\!E \left[\|U_{n,\alpha_n}(t)u_n^0 - p_n \exp(tA)u^0\| \right] \le C\alpha_n \|A \exp(tA)u^0\| + \|u_{n,\alpha_n}(t) - p_n u_{\alpha_n}(t)\| + C\tau_n \|u_n^0\|, \quad 0 < t \le T.$$

5 Coercive inequalities

Let us consider in the Banach space E the nonhomogeneous Cauchy problem:

$$\begin{aligned} &u'(t) = Au(t) + f(t),\ t \in [0,T], \\ &u(0) = u_0, \end{aligned} \tag{5.1}$$

with operator A which generates C_0-semigroup and $f(\cdot)$ be some function from $[0,T]$ to E. The problem (5.1) can be considered in different functional spaces. The most popular situations are the following settings: well-posedness in $C([0,T];E), C^{\alpha,0}([0,T];E)$ and $L^p([0,T];E)$ spaces (see [9], [22], [136], [214]).

We say that problem (5.1) is well-posed, say in $C([0,T];E)$, if for any $f(\cdot) \in C([0,T];E)$ and any $u_0 \in D(A)$ the problem (5.1) is

i). uniquely solvable, i.e. $u(\cdot)$ satisfies the equation and boundary condition (5.1) , $u(\cdot)$ is continuously differentiable on $[0,T]$, $u(t) \in D(A)$ for any $t \in [0,T]$ and $Au(\cdot)$ is continuous on $[0,T]$;

ii). operator $(f(\cdot), u^0) \to u(\cdot)$, regarded as an operator from $C([0,T];E) \times D(A)$ to $C([0,T];E)$ is continuous.

In case $u^0 \equiv 0$, the coercive well-posedness in $C([0,T];E)$ means that $\|Au(\cdot)\|_{C([0,T];E)} \le c\|f(\cdot)\|_{C([0,T];E)}$. In general coercive well-posedness in the space $\Upsilon([0,T];E)$ means for problem (5.1) that it is well-posed in the space $\Upsilon([0,T];E)$ and one has

$$\|u'(\cdot)\|_{\Upsilon([0,T];E)} + \|Au(\cdot)\|_{\Upsilon([0,T];E)} \le C\left(\|f(\cdot)\|_{\Upsilon([0,T];E)} + \|u^0\|_F\right),$$

where F is some subspace of E. For results of coercive well-posedness see [9], [22], [136].

The semidiscrete approximation of (5.1) are the Cauchy problems in Banach spaces E_n :

$$\begin{aligned} &u_n'(t) = A_n u_n(t) + f_n(t),\ t \in [0,T], \\ &u_n(0) = u_n^0, \end{aligned} \tag{5.2}$$

with operators A_n which generate C_0-semigroups, A_n and A are compatible, $u_n^0 \to u^0$ and $f_n \to f$ in appropriate sense. Following section 3 it is natural to assume that conditions (A) and (B_1) are satisfied.

Here we are going to describe the discretization of (5.2) in time. The simplest difference scheme (Rothe scheme) is

$$\begin{array}{ll} \frac{\overline{U}_n^k - \overline{U}_n^{k-1}}{\tau_n} & = A_n \overline{U}_n^k + \varphi_n^k,\ k \in \{1, ..., [\frac{T}{\tau_n}]\}, \\ \overline{U}_n^0 & = u_n^0, \end{array} \tag{5.3}$$

where, for example, in the case of $f_n(\cdot) \in C([0,T]; E_n)$ one can put $\varphi_n^k = f_n(k\tau_n), k \in \{1, ..., K\}, K = [\frac{T}{\tau_n}]$, and in the case $f_n \in L^1([0,T]; E_n)$ one can set $\varphi_n^k = \frac{1}{\tau_n} \int_{t_{k-1}}^{t_k} f_n(s) ds,\ t_k = k\tau_n, k \in \{1, ..., K\}$.

5.1 Coercive inequality in $C_{\tau_n}([0,T]; E_n)$ spaces

Denote by $C_{\tau_n}([0,T]; E_n)$ the space of the elements $\overline{\varphi}_n = \{\varphi_n^k\}_{k=0}^K$ such that $\varphi_n^k \in E_n, k \in \{0, ..., K\}$, with the norm $\|\overline{\varphi}_n\|_{C_{\tau_n}([0,T];E_n)} = \max_{0 \le k \le K} \|\varphi_n^k\|_{E_n}$.

We recall that coercive well-posedness in $C([0,T]; E)$ implies [22] that A generates an analytic C_0-semigroup.

Theorem 5.1 *[22] Let condition (B_1) be satisfied. The problem (5.3) is stable in the space $C_{\tau_n}([0,T]; E_n)$, i.e.*

$$\|\overline{U}_n\|_{C_{\tau_n}([0,T];E_n)} \le C\Big(\|\overline{\varphi}_n\|_{C_{\tau_n}([0,T];E_n)} + \|u_n^0\|\Big).$$

Theorem 5.2 *[22] Let condition (B_1) be satisfied. The problem (5.3) is almost coercive stable in the space $C_{\tau_n}([0,T]; E_n)$, i.e.*

$$\|A_n \overline{U}_n\|_{C_{\tau_n}([0,T];E_n)} \le M\Big(\|A_n u_n^0\|_{E_n} + \min\Big(\ln(1/\tau_n), 1 + \Big|\ln \|A_n\|\Big|\Big)\|\overline{\varphi}_n\|_{C_{\tau_n}([0,T];E_n)}\Big).$$

We have note that if (5.1) is coercive well-posed in the space $C([0,T]; E)$, then [66] operator A should be bounded or the space E contains a subspace isomorphic to c_0. It means that in general problem (5.3) is not coercive well-posed in $C_{\tau_n}([0,T]; E_n)$ space.

For the explicit scheme

$$\begin{array}{ll} \frac{U_n^k - U_n^{k-1}}{\tau} & = A_n U_n^{k-1} + \varphi_n^k,\ k \in \{1, ..., K\}, \\ U_n^0 & = u_n^0, \end{array} \tag{5.4}$$

Theorem 5.2 could be reconstructed, but under a stability condition.

Theorem 5.3 *[22] Let condition (B_1) is satisfied and $\tau_n \ln(\frac{1}{\tau_n}) \|A_n\| \le \epsilon$ for sufficiently small $\epsilon > 0$. Then the problem (5.4) is almost coercive stable in the space $C_{\tau_n}([0,T]; E_n)$, i.e.*

$$\begin{aligned} &\|A_n U_n\|_{C_{\tau_n}([0,T];E_n)} + \|U_n\|_{C_{\tau_n}([0,T];E_{n,1-\frac{1}{\ln \frac{1}{\tau_n}}})} \le \\ \le\ & M\Big(\|A_n u_n^0\|_{E_{n,1-\frac{1}{\ln \frac{1}{\tau_n}}}} + \min\Big(\ln(1/\tau_n), 1 + \Big|\ln \|A_n\|\Big|\Big)\|\overline{\varphi}_n\|_{C^{\tau_n}([0,T];E_n)}\Big), \end{aligned}$$

where $\|u_n\|_{E_{n,\alpha}} = \Big(\int_0^\infty \|A_n \exp(tA_n) u_n\|_{E_n}^{\frac{1}{1-\alpha}} dt\Big)^{1-\alpha}$.

Remark 5.1 *The space $E_{n,\alpha}$ coincides with equivalent norm with the real interpolation space $(E_n, D(A_n))_{1-1/p,p}$ see [136].*

5.2 Coercive inequality in $C^{\alpha,0}_{\tau_n}([0,T];E_n)$ spaces

Denote by $C^{\alpha,0}_{\tau_n}([0,T];E_n)$ for $0<\alpha<1$ the space of the elements $\overline{\varphi}_n$ with the norm $\|\overline{\varphi}_n\|_{C^{\alpha,0}_{\tau_n}([0,T];E_n)} = \max_{0\le k\le K}\|\varphi_n^k\|_{E_n} + \max_{1\le k<k+l\le K}\|\varphi_n^{k+l}-\varphi_n^k\|_{E_n}(\tau_n k)^\alpha (l\tau_n)^{-\alpha}$.

Theorem 5.4 *[180] Let condition (B_1) hold. Then the scheme (5.3) is coercive well-posed in $C^{\alpha,0}_{\tau_n}([0,T];E_n)$ with $0<\alpha<1$, i.e.*

$$\|A_n\overline{U}_n\|_{C^{\alpha,0}_{\tau_n}([0,T];E_n)} \le \frac{M}{\alpha(1-\alpha)}\Big(\|A_n u_n^0\|_{E_n} + \|\overline{\varphi}_n\|_{C^{\alpha,0}_{\tau_n}([0,T];E_n)}\Big).$$

Roughly speaking, assumption (B_1) is necessary and sufficient for coercive well-posedness in $C^{\alpha,0}_{\tau_n}([0,T];E_n)$ space.

5.3 Coercive inequality in $L^p_{\tau_n}([0,T];E_n)$ spaces

Denote by $L^p_{\tau_n}([0,T];E_n)$ for $1\le p<\infty$ the space of the elements $\overline{\varphi}_n$ with the norm $\|\overline{\varphi}_n\|_{L^p_{\tau_n}([0,T];E_n)} = \Big(\Sigma_{j=0}^K\|\varphi_n^k\|^p_{E_n}\tau_n\Big)^{1/p}$.

Theorem 5.5 *[180] Let condition (B_1) hold. Let the difference scheme (5.3) be coercive well-posed in $L^{p_0}_{\tau_n}([0,T];E_n)$ for some $1<p_0<\infty$. Then it is coercive well-posed in $L^p_{\tau_n}([0,T];E_n)$ for any $1<p<\infty$ and*

$$\|A_n\overline{U}_n\|_{L^p_{\tau_n}([0,T];E_n)} + \max_{0\le k\le K}\|\overline{U}_n^k\|_{E_{n,1-1/p}} \le \frac{Mp^2}{p-1}\Big(\|\overline{\varphi}_n\|_{L^p_{\tau_n}([0,T];E_n)} + \|\overline{U}_n^0\|_{1-1/p}\Big).$$

We have to note that to the contrary to $C^{\alpha,0}$ space case the analyticity of semigroup $\exp(\cdot A)$ is not enough to get coersive well-posedness in L^p space [124], so to state coercive well-posedness in L^p we need some additional assumptions.

Theorem 5.6 *[180] Let $1<p,q<\infty, 0<\alpha<1$ and condition (B_1) hold. Then the difference scheme (5.3) is coercive well-posed in $L^p_{\tau_n}([0,T];E_{n,\alpha,q})$, i.e.*

$$\|A_n\overline{U}_n\|_{L^p_{\tau_n}([0,T];E_{n,\alpha,q})} + \max_{0\le k\le K}\|U_n^k\|_{E_{n,1-1/p}} \le \frac{Mp^2}{(p-1)\alpha(1-\alpha)}\Big(\|\overline{\varphi}_n\|_{L^p_{\tau_n}([0,T];E_{n,\alpha,q})} + \|U_n^0\|_{1-1/p}\Big),$$

where $E_{n,\alpha,q}$ is the interpolation space $(E_n, D(A_n))_{\alpha,q}$ with the norm $\|u_n\|_{E_{n,\alpha,q}} = \Big(\int_0^\infty\|\lambda^\alpha A_n(\lambda - A_n)^{-1}\|^q_{E_n}\frac{d\lambda}{\lambda}\Big)^{1/q}$.

For general Banach space E we have the following results. Assume that A is the generator of the analytic semigroup $\exp(tA), t\in I\!R_+$, of the linear bounded operators with exponentially decreasing norm when $t\to\infty$. That means that stability condition (B_1'') holds with $\omega''\le 0$.

Theorem 5.7 *[21] Let condition (B_1) hold. Then the solution of difference scheme (5.3) is almost coercive stable, i.e.*

$$\|A_n\overline{U}_n\|_{L^p_{\tau_n}([0,T];E_n)} \le M\Big(\|A_n\overline{U}_n^0\|_{E_n} + \min\{\ln\frac{1}{\tau_n}, 1+|\ln\|A_n\|_{B(E_n)}|\}\,\|\varphi_n\|_{L^p_{\tau_n}([0,T];E_n)}\Big)$$

holds for any $p\ge 1$, where M does not depend on τ_n, u_n^0, and φ_n.

Of course for schemes like

$$\begin{array}{ll} \frac{U_n^k - U_n^{k-1}}{\tau_n} & = A_n(\frac{U_n^k + U_n^{k-1}}{2}) + \varphi_n^k,\ n \in \{1, ..., K\}, \\ U_n^0 & = u_n^0. \end{array} \tag{5.5}$$

coercive well-posedness could be considered.

Theorem 5.8 *[21] Let the condition (B_1) holds. Then the solution of difference scheme (5.5) is almost coercive stable, i.e. the estimate*

$$\|\{A_n \frac{U_n^j + U_n^{j-1}}{2}\}\|_{L^p_{\tau_n}([0,T];E_n)} \le M\Big(\|A_n u_n^0\|_{E_n} + \min\{\ln \frac{1}{\tau_n},\ 1 + |\ln \|A_n\|_{E_n \mapsto E_n}|\}\ \|\varphi_n\|_{L^p_{\tau_n}([0,T];E_n)}\Big)$$

holds for any $p \ge 1$, where M does not depend on τ_n, u_n^0, and φ_n.

Theorem 5.9 *[21] Let the condition (B_1) holds and condition (3.6) is satisfied. Then the solution of difference scheme (5.5) is almost coercive stable, i.e. the estimate*

$$\|A_n U_n\|_{L^p_{\tau_n}([0,T];E_n)} \le M\Big(\|A_n u_n^0\|_{E_n} + \min\{\ln \frac{1}{\tau_n},\ 1 + |\ln \|A_n\|_{E_n \mapsto E_n}|\}\ \|\varphi_n\|_{L^p_{\tau_n}([0,T];E_n)}\Big)$$

holds for any $p \ge 1$, where M does not depend on τ_n, u_n^0, and φ_n.

The necessary and sufficient conditions for coercive well-posedness of the problem (5.1) in $L^p([0,T];E)$ was obtained in [219], [218], [98]. More precisely, a Banach space E has the UMD–property, whenever the Hilbert transform $Hf(t) = \frac{1}{\pi} PV - \int_{-\infty}^{\infty} \frac{1}{t-s} f(s) ds$ extends to a bounded operator on $L^p(I\!R; E)$ for some (all) $p \in (1, \infty)$. It is well known, that all subspaces and quotient spaces of $L^q(\Omega, \mu)$ with $1 < q < \infty$ have this property.

The Poisson semigroup on $L^1(I\!R)$ and on $L^p(I\!R; E)$ does not coercive well posed on $L^p(I\!R, E)$ space if E is not an UMD–space (cf. [124]). Hence the assumptions on E to be an UMD space is necessary in some sense.

But it was an open problem whether every generator of an analytic semigroup on $L^q(\Omega, \mu)$, $1 < q < \infty$, provide coercive well-posedness in $L^p(I\!R; E)$. Recently Kalton and Lancien [100] gave a strong negative answer to this question. If every bounded analytic semigroup on a Banach space E is such that problem (5.1) is coercive well-posed, then E is isomorphic to a Hilbert space.

If A generates a bounded analytic semigroup $\{\exp(zA) : |\arg(z)| \le \delta\}$, on a Banach space E, then the following three sets are bounded in the operator norm

i) $\{\lambda(\lambda - A)^{-1} : \lambda \in iI\!R, \lambda \ne 0\}$;

ii) $\{\exp(tA), tA\exp(tA) : t > 0\}$;

iii) $\{\exp(zA) : |\arg z| \le \delta\}$.

In Hilbert spaces this already implies coercive well-posedness in $L^p(I\!R_+; E)$, but only in Hilbert spaces E. The additional assumption that we need in more general Banach spaces E will be R–boundedness.

A set $\mathcal{T} \subset B(E)$ is called R–bounded, if there is a constant $C < \infty$, such that for all $Z_1, \ldots, Z_k \in \mathcal{T}$ and $x_1, \ldots, x_k \in E, k \in I\!N$,

$$\int_0^1 \|\sum_{j=0}^k r_j(u) Z_j(x_j)\|\, du \le C \int_0^1 \|\sum_{j=0}^k r_j(u) x_j\|\, du, \tag{5.6}$$

where $\{r_j\}$ is a sequence of independent symmetric $\{-1, 1\}$– valued random variables, e. g. the Rademacher functions $r_j(t) = \text{ sign } (\sin(2^j \pi t))$ on $[0, 1]$. The smallest C, such that (5.6) is fulfilled, is called the R–boundedness constant of $\mathcal{T}$ and is denoted by $R(\mathcal{T})$.

Theorem 5.10 *[219] Let A generates a bounded analytic semigroup $\exp(tA)$ on a UMD–space E. Then the problem (5.1) is coercive well-posed in the space $L^p(I\!R_+;E)$ if and only if one of the sets i), ii) or iii) above is R–bounded.*

The interpretation of discrete coercive inequality and discrete semigroup define the convolution operator in the form $\breve{A}_n\Sigma_{j=0}^k T_n^{k-j}Q_n\varphi_n\tau_n$ with some bounded operator $Q_n \in B(E_n)$, which usually has smoothness property as it is clear from the Proofs of the Theorems 5.7 and 5.8. Here $T_n(\tau_n)^k$ is discrete semigroup say as in Subsection 3.1. Boundedeness of convolution operator in $L^p_{\tau_n}(Z\!\!Z_+;E_n)$ space implies discrete coercive well-posedness in $L^p_{\tau_n}(Z\!\!Z_+;E_n)$.

We assume also that Banach spaces E_n satisfy in this section the collectively UMD–property, i.e. we assume that the Hilbert transforms $H_nf_n(t) = \frac{1}{\pi}PV - \int_{-\infty}^{\infty}\frac{1}{t-s}f_n(s)ds$ extend to a bounded operators on $L^p(I\!R;E_n)$ for some (all) $p \in (1,\infty)$, such that all of them are bounded by constant which does not depend on n. This assumption holds for example if all E_n can be embedded into a fixed space $L^p(\Omega)$ with $1 < p < \infty$.

Definition 5.1 *A discrete semigroup $T_n(\cdot)$ with generator $\breve{A}_n$ is said generates coercive well-posedness on $L^p_{\tau_n}(Z\!\!Z_+;E_n)$ space if the corresponding convolution operator $\varphi_n \mapsto \{\breve{A}_n\Sigma_{j=0}^k T_n^{k-j}Q_n\varphi_n^j\tau_n\}$ is continuous on $L^p_{\tau_n}(Z\!\!Z_+;E_n)$ space.*

Theorem 5.11 *[21] Assume that for convolution operator $\varphi_n \mapsto \{\breve{A}_n\Sigma_{j=0}^k T_n^{k-j}Q_n\varphi_n^j\tau_n\}$ the following conditions hold*

1^0. the set $\{\breve{A}_n(\lambda - T_n)^{-1}Q_n\tau_n : |\lambda| = 1, \lambda \neq 1, \lambda \neq -1\}$ is R-bounded;

2^0. the set $\{(\lambda-1)(\lambda+1)\breve{A}_n(\lambda - T_n)^{-2}Q_n\tau_n : |\lambda| = 1, \lambda \neq 1, \lambda \neq -1\}$ is R-bounded.

Then a discrete semigroup $T_n(\cdot)$ generates coercive well-posedness on $L^p_{\tau_n}(Z\!\!Z_+;E_n)$ space.

Theorem 5.12 *[21] Let E_n be UMD Banach spaces. Assume also that the set $\{\lambda(\lambda - A_n)^{-1} : \lambda \in i I\!R, \lambda \neq 0\}$ is R-bounded with the R–boundedness constant which does not depend on n. Then the solution of difference scheme (5.3) is coercive stable, i.e.*

$$\|\breve{A}_n\overline{U}_n\|_{L^p_{\tau_n}(Z\!\!Z_+;E_n)} \leq M\|\varphi_n\|_{L^p_{\tau_n}(Z\!\!Z_+;E_n)} \tag{5.7}$$

holds for any $p \geq 1$, where M does not depend on τ_n, u_n^0, and φ_n.

Remark 5.2 *We have to note that Theorem 3.2 could be reformulates in the notions of R-boundedness with changes of condition (B_1) for condition: there is a $0 < \theta < \pi/2$ such that the set $\{\lambda(\lambda - A_n)^{-1} : \lambda \in \Sigma(\theta + \pi/2)\}$ is R-bounded with the R–boundedness constant which does not depend on n. The condition (C_1) could be written, due to [219] Theorem 4.2, in the form $\exp(tA_n) \to \exp(tA)$ converges for any $t \in I\!R$ and there is a $0 < \theta < \pi/2$ such that the set $\{\exp(zA_n) : z \in \Sigma(\theta)\}$ is R-bounded with the R–boundedness constant which does not depend on n. So one of our assumption in Theorems 5.12 and 5.13 is in some sense condition (B_1) changed for R-boundedness condition.*

Theorem 5.13 *[21] Let E_n be UMD Banach spaces. Assume also that the set $\{\lambda(\lambda - A)^{-1} : \lambda \in i I\!R, \lambda \neq 0\}$ is R-bounded with the R–boundedness constant which does not depend on n. Then the solution of difference scheme (5.5) is coercive stable, i.e.*

$$\|\{\breve{A}_n\frac{U_n^k + U_n^{k-1}}{2}\}\|_{L^p_{\tau_n}([0,T];E_n)} \leq M\|\varphi_n\|_{L^p_{\tau_n}([0,T];E_n)} \tag{5.8}$$

holds for any $p \geq 1$, where M does not depend on τ_n, u_n^0, and φ_n.

Remark 5.3 *Analyzing the Proofs of Theorems 5.12 and 5.13 it is easy to see that one can put* $\breve{A}_n = A_n$ *in the statements (5.7) and (5.8). Moreover, the statement (5.8) could be written in the form*

$$\|\breve{A}_n U_n\|_{L^p_{\tau_n}([0,T];E_n)} \le M\|\varphi_n\|_{L^p_{\tau_n}([0,T];E_n)}.$$

The proof of that fact based on equality $\breve{A}_n = A_n(I_n - \frac{\tau_n}{2}A_n)^{-1}$.

It is possible to consider a more general Padé difference scheme [22] for $p = q-1$ or $p = q-2$. In this case the difference scheme will be written in the form

$$\frac{U_n^k - U_n^{k-1}}{\tau_n} = (\breve{A}_n U_n)^{k-1} + \varphi_n^{p,q,k},\ U_n^0 = u_n^0,\ 1 \le k \le K. \tag{5.9}$$

where $(\breve{A}_n U_n)^k = (\frac{R_{p,q}(\tau_n A_n)-I}{\tau_n}U_n)^{k-1}$ and $\|\varphi_n^k - \varphi_n^{p,q,k}\|_{E_n} \le M\tau_n^{p+q}$. To formulate coercive statements of subsections 5.1-5.3 we just need to change operator A_n for $\breve{A}_n$. We know from Theorem 3.16 that Padé approximation under condition (B_1) with $p = q$ is stable, but in general it will be not coercive stable. To get coercive inequality we need condition (3.6). Spaces where the problem considered also could be very different [22].

5.4 Coercive inequality in $B_{\tau_n}([0,T];C_h^\theta(\Omega_h)) \cap C_h([0,T];C_h(\bar{\Omega}_h))$

From the numerical analysis point of view it would be very interesting to consider the Problem (5.1) in the space $\Upsilon([0,T];E)$ that E would be smoother than $C(\Omega)$ (elements of such a space could easily be well approximated) and $\Upsilon([0,T];E)$ be like $C([0,T];E)$ or bounded functions space. The interesting fact is that such a situation actually possible at least for the strongly elliptic operator of order 2 with coefficients of class $C^\theta(\overline{\Omega})$. Since in such space E operator $(p_n v)_i = v(ih)$ very concrete, i.e. taking values in the grid points, we consider the case omitting in this section the notion of p_n.

Theorem 5.14 *[36] Let* Ω *be an open bounded subset of* $\mathbb{R}^d$*, lying on one side of its topological boundary* $\partial\Omega$*, which is a submanifold of* $\mathbb{R}^d$ *of dimension* $d-1$ *and class* $C^{2+\theta}$*, for some* $\theta \in (0,2)\setminus\{1\}$*. Let* $\mathcal{A} = \mathcal{A}(x,D_x) = \sum_{|\alpha|\le 2} a_\alpha(x)D_x^\alpha$ *be a strongly elliptic operator of order two (thus,* $Re\sum_{|\alpha|=2} a_\alpha(x)\xi^\alpha \ge \nu|\xi|^2$ *for some* $\nu > 0$ *and for any* $(x,\xi) \in \overline{\Omega} \times \mathbb{R}^d$*) with coefficients of class* $C^\theta(\overline{\Omega})$*. Then, there exist* $\mu \ge 0$*,* $\phi_0 \in (\frac{\pi}{2},\pi)$ *such that for any* $\lambda \in \mathbb{C}$*, with* $|\lambda| \ge \mu$ *and* $|Arg\lambda| \le \phi_0$ *the problem*

$$\begin{aligned} &\lambda v - \mathcal{A}v = y,\\ &\gamma_0 v = 0, \end{aligned}$$

has for any $y \in C^\theta(\overline{\Omega})$ *a unique solution* v *belonging to* $C^{2+\theta}(\overline{\Omega})$ *and for a certain* $M > 0$*,*

$$|\lambda|^{1+\frac{\theta}{2}}\|v\|_{C(\overline{\Omega})} + |\lambda|\,\|v\|_{C^\theta(\overline{\Omega})} + \|v\|_{C^{2+\theta}(\overline{\Omega})} \le M\Big(\|y\|_{C^\theta(\overline{\Omega})} + |\lambda|^{\frac{\theta}{2}}\|\gamma_0 y\|_{C(\partial\Omega)}\Big), \tag{5.10}$$

where γ_0 *is the trace operator on* $\partial\Omega$*.*

It is clear from (5.10) that operator $\mathcal{A}$ in general does not generate C_0-semigroup in $E = C^\theta(\overline{\Omega})$ space, but, following say [136], one can construct semigroup $\exp(t\mathcal{A}), t \ge 0$, which will be analytic.

Let $\mathcal{I} = \mathbb{Z}$ and E be a Banach space with norm $\|\cdot\|$. For a grid function $U : \mathcal{I} \to E$, writing U_j or $(U)_j$ instead of $U(j)$ for any $j \in \mathcal{I}$, we set

$$B(\mathcal{I};E) := \{U : \mathcal{I} \to E \ :\ \sup_{j\in\mathcal{I}}\|U_j\| < +\infty\},\ \|U\|_{B(\mathcal{I};E)} := \sup_{j\in\mathcal{I}}\|U_j\|.$$

It is easily seen that $B(\mathcal{I};E)$ is a Banach space with the norm $\|\cdot\|_{B(\mathcal{I};E)}$. If the set $\mathcal{I}$ is some interval, say $\mathcal{I}=(a,\infty)$, we denote by $B(\mathcal{I};E)$ the set of all bounded functions from $\mathcal{I}$ to E.

For the grid function $U:\mathcal{I}\to E$ and $h>0$, we define operator ∂_h by formula

$$(\partial_h U)_j := h^{-1}(U_{j+1}-U_j).$$

For any $m\in\mathbb{Z}$ we set $(\partial_h^m U)_j := h^{-m}\sum_{i=0}^m \binom{m}{i}(-1)^{m-i}U_{j+i}$. If $U\in B(\mathcal{I};E)$, we set

$$\|U\|_{C_h^m(\mathcal{I};E)} := \max\Big\{\|\partial_h^r U\|_{B(\mathcal{I};E)} \,:\, 0\le r\le m\Big\}.$$

Finally, let $\theta\in(0,1)$. We define

$$[U]_{C_h^\theta(\mathcal{I};E)} := \sup\Big\{\Big((k-j)h\Big)^{-\theta}\|U_k-U_j\| \,:\, j,k\in\mathcal{I}, j<k\Big\}$$

and, if $m\in I\!N_0$,

$$\|U\|_{C_h^{m+\theta}(\mathcal{I};E)} := \max\Big\{\|U\|_{C_h^m(\mathcal{I};E)}, [\partial_h^m U]_{C_h^\theta(\mathcal{I},E)}\Big\}.$$

In the same context we shall indicate $B(\mathcal{I};E)$ with $C_h^0(\mathcal{I};E)$. If $E=\mathbb{C}$ we write simply $B(\mathcal{I})$ or $C_h^0(\mathcal{I})$.

Let $f\in B(I\!N;E)$. We indicate with $\tilde f$, the extension of f to $I\!N_0$ such that $\tilde f_0=0$. For a nonnegative real number ω, we define

$$\|f\|_{C^\omega_{h,0}(I\!N;E)} := \|\tilde f\|_{C^\omega_h(I\!N_0;E)}. \tag{5.11}$$

Let now $L>0, n\in I\!N, n\ge 3$, and $h=\frac{L}{n}$. For $j\in\mathcal{I}:=\{1,...,n-1\}$ we are given complex numbers a_j,b_j,b'_j,c_j satisfying the following conditions (ι):

$(\iota 1)$: there exists $\nu>0$ such that $Re(a_j)\ge\nu$, for every $j\in\mathcal{I}$;

$(\iota 2)$: $\max\Big\{|a_j|,|b_j|,|b'_j|,|c_j|\Big\}\le Q$ for every $j\in\mathcal{I}$ with $Q>\nu$;

$(\iota 3)$; there exists $\omega:[0,L]\to[0,+\infty)$ such that $\omega(0)=0$ and ω is continuous in 0 such that for $j,k\in\mathcal{I}$ with $j\le k$

$$|a_k-a_j|\le\omega\Big((k-j)h\Big).$$

For $\lambda\in\mathbb{C}$ we are going to study the following problem:

$$\begin{array}{l}\lambda U_j - a_j(\partial_h^2 U)_{j-1} - b_j(\partial_h U)_j - b'_j(\partial_h U)_{j-1} - c_j U_j = f_j \quad \text{for } j=1,...,n-1,\\ U_0=U_n=0.\end{array} \tag{5.12}$$

To this aim, we set $\overline{\mathcal{I}}:=\{0,1,...,n-1,n\}$ and for $U\in B(\mathcal{I};E)$, we define

$$\tilde U_j = \begin{cases} U_j & \text{if } j\in\mathcal{I},\\ 0 & \text{if } j\in\{0,n\}.\end{cases}$$

We introduce operator A_h in $B(\mathcal{I};E)$ defined as follows:

$$(A_hU)_j := a_j(\partial_h^2\tilde U)_{j-1} + b_j(\partial_h\tilde U)_j + b'_j(\partial_h\tilde U)_{j-1} + c_j\tilde U_j \ \text{ for } j\in\mathcal{I}. \tag{5.13}$$

Further, we assume that

$(\iota_\theta 1)$: there exists $\nu>0$ such that $Re(a_j)\ge\nu$ for every $j\in\overline{\mathcal{I}}$;

$(\iota_\theta 2)$: $\max\Big\{\|a\|_{C_h^\theta(\overline{\mathcal{I}})}, \|b\|_{C_h^\theta(\overline{\mathcal{I}})}, \|b'\|_{C_h^\theta(\overline{\mathcal{I}})}, \|b\|_{C_h^\theta(\overline{\mathcal{I}})}\Big\}\le Q$ with $Q>\nu$.

Proposition 5.1 *[92] Assume that assumptions (ι_θ) are satisfied for some $\theta \in (0,2)\setminus\{1\}$. Fix $\phi_0 \in [0, \pi - \arccos(\frac{\nu}{Q}))$. Then there exists $\mu_0 > 0$ such that $\{\lambda \in \mathbb{C} : |\lambda| \geq \mu_0, |Arg(\lambda)| \leq \phi_0\} \subseteq \rho(A_h)$ where A_h is the operator defined in (5.13). Moreover, for every $r \in [0,2]$ there exists $c > 0$ depending only on L, ν, Q, r such that for every $f \in B(\mathcal{I}; E)$, and any $F \in B(\overline{\mathcal{I}}; E)$ with $F|_{\mathcal{I}} = f$, one has*

$$\|(\lambda - A_h)^{-1} f\|_{C_{h,0}^{\theta+r}(\mathcal{I};E)} \leq c|\lambda|^{\frac{r}{2}-1}\Big(\|F\|_{C_h^\theta(\overline{\mathcal{I}};E)} + |\lambda|^{\frac{\theta}{2}} \max\{\|F_0\|, \|F_n\|\}\Big).$$

Let us consider the following mixed Cauchy-Dirichlet parabolic problem:

$$\begin{array}{ll} \frac{\partial u}{\partial t}(t,x) = \mathcal{A}u(t,x) + f(t,x), & t \in [0,T], x \in [0,L], \\ u(t,x') = 0, & t \in [0,T], x' \in \{0,L\}, \\ u(0,x) = 0, & x \in [0,L], \end{array} \tag{5.14}$$

where $\mathcal{A}$ is a second order differential operator and $L > 0$. We say that problem (5.14) has a strict solution if there exists a continuous function $u(t,x)$ with the first derivative with respect to t and the derivatives of order less than or equal to 2 with respect to x which are continuous up to boundary of $[0,T] \times [0,L]$, i.e. $u \in C^1\Big([0,T]; C(\overline{\Omega})\Big) \cap C\Big([0,T]; C^2(\overline{\Omega})\Big)$, and the equations in (5.14) are satisfied.

Theorem 5.15 *[88] Consider problem (5.14) under the following assumptions:*

(I) T and L are positive real numbers;

(II) $\theta \in (0,1)\setminus\{\frac{1}{2}\}$;

(III) $\mathcal{A}u(x) = a(x)\frac{\partial^2 u}{\partial x^2}(t,x) + b(x)\frac{\partial u}{\partial x}(t,x) + c(x)u(t,x)$, with $a, b, c \in C^{2\theta}([0,L])$;

(IV) a is real valued and $\min a = \nu > 0$;

(V) $f \in C([0,T]\times[0,L])$, $t \to f(t,.) \in B\Big([0,T]; C^{2\theta}([0,L])\Big)$, $t \to f(t,0)$ and $t \to f(t,L)$ belong to $C^\theta([0,T])$, $f(0,0) = f(0,L) = 0$.

Then there exists a unique strict solution $u(\cdot)$ of problem (5.14). Such solution belongs to $B\Big([0,T]; C^{2+2\theta}([0,L])\Big)$ and $\frac{\partial u}{\partial t} \in B\Big([0,T]; C^{2\theta}([0,L]\Big)$.

Now let $\mathcal{I}$ be a set which can depend on a positive parameter h and the Banach space $X_h = B(\mathcal{I})$. Next we introduce a linear operator A_h in X_h depending on h. In every case $\rho(A_h)$ contains $\{\lambda \in \mathbb{C}\setminus\{0\} : |\lambda| \geq R \text{ and } |Arg(\lambda)| \leq \phi_0\}$, where $R > 0$ and $\phi_0 \in (\frac{\pi}{2}, \pi)$, and there exists $M > 0$ such that for λ in the specified subset of $\mathbb{C}$, $\|(\lambda - A_h)^{-1}\|_{L(X_h)} \leq M|\lambda|^{-1}$. Here R, ϕ_0 and M are independent of h. Then we consider another set $\tilde{\mathcal{I}}$ such that $\mathcal{I} \subseteq \tilde{\mathcal{I}}$, we put $\tilde{X}_h := B(\tilde{\mathcal{I}})$. We define an extension operator $\mathscr{E}_h$ from X_h to $\tilde{X}_h$: in all our concrete cases this was the extension with zero. Next, for $\theta \in (0,1)$, we introduce the norms $\|\cdot\|_{2\theta,h}$ and $\|\cdot\|_{2+2\theta,h}$ in $\tilde{X}_h$. The first of these norms is connected to $\|\cdot\|_X$ and the operator A_h by the following property: there exist two positive constants c_1 and c_2 independent of h such that for every $U \in X_h$

$$c_1\|\mathscr{E}_h U\|_{2\theta,h} \leq \|U\|_{(X_h, D(A_h))_\theta} \leq c_2\|\mathscr{E}_h U\|_{2\theta,h}.$$

Then for every h we consider a restriction operator $\mathcal{R}_h \in L(\tilde{X}_h, X_h)$, such that $\mathcal{R}_h\mathscr{E}_h = I_{X_h}$. Let us also introduce a seminorm p_h in $\tilde{X}_h$: in the concrete cases we have $p_h(U) = \|U|_{\tilde{\mathcal{I}}\setminus\mathcal{I}}\|_{B(\tilde{\mathcal{I}}\setminus\mathcal{I})}$. We assume that, if $|\lambda| \geq R$ and $|Arg(\lambda)| \leq \phi_0$, for every $G \in \tilde{X}_h$, then

$$|\lambda|\,\|\mathscr{E}_h(\lambda - A_h)^{-1}\mathcal{R}_h G\|_{2\theta,h} + \|\mathscr{E}_h(\lambda - A_h)^{-1}\mathcal{R}_h G\|_{2+2\theta,h} \leq M\Big(\|G\|_{2\theta,h} + |\lambda|^\theta p_h(G)\Big).$$

Another inequality we impose is the following. If $|\lambda| \geq R$, $|Arg(\lambda)| \leq \phi_0$ and $G \in \tilde{X}_h$, then

$$\|A_h(\lambda - A_h)^{-1}\mathcal{R}_h G\|_{X_h} \leq M|\lambda|^{-\theta}\Big(\|G\|_{2\theta,h} + |\lambda|^\theta p_h(G)\Big).$$

Such an inequality can be easily deduced in each of our examples. In the formulation below we remove the parameter h. In the case of the backward Euler scheme (5.3) we have

Theorem 5.16 *[92] Let X and $\tilde{X}$ be Banach spaces with norms $\|\cdot\|_X$ and $\|\cdot\|_{\tilde{X}}$ respectively, $A \in B(X), \mathscr{E} \in B(X,\tilde{X}), \mathcal{R} \in B(\tilde{X},X)$ such that $\mathcal{R}\mathscr{E} = I_X$. Assume, moreover, that $\theta \in (0,1)$ and $\|\cdot\|_{2\theta}$ and $\|\cdot\|_{2+2\theta}$ are norms in $\tilde{X}$, while p is a seminorm in the same space. Finally, assume that there exist $R > 0, \phi_0 \in (\frac{\pi}{2}, \pi), M > 0$, such that the following conditions are satisfied:*

(a) $\{\lambda \in \mathbb{C} : |\lambda| \geq R, |Arg(\lambda)| \leq \phi_0\} \subseteq \rho(A)$ and, for λ in this set,

$$\|(\lambda - A)^{-1}\|_{B(X)} \leq M(1 + |\lambda|)^{-1};$$

(b) for every $F \in X$

$$\|\mathscr{E}F\|_{2\theta} \leq M\|F\|_{(X,D(A))_\theta};$$

(c) for every $V \in \tilde{X}, \lambda \in \mathbb{C}$ with $|\lambda| \geq R$ and $|Arg(\lambda)| \leq \phi_0$

$$(1 + |\lambda|)^{-1}\|\mathscr{E}(\lambda - A)^{-1}\mathcal{R}V\|_{2\theta} + \|\mathscr{E}(\lambda - A)^{-1}\mathcal{R}V\|_{2+2\theta}$$

$$+(1 + |\lambda|)^{\theta}\|A(\lambda - A)^{-1}\mathcal{R}V\|_X \leq M\Big(\|V\|_{2\theta} + (1 + |\lambda|)^{\theta}p(V)\Big);$$

(d) $p(V) \leq \|V\|_{2\theta}$ for every $V \in \tilde{X}$ and $p(\mathscr{E}F) = 0$ for every $F \in X$;

(e) $\|\mathcal{R}V\|_X \leq \|V\|_{2\theta}$ for every $V \in \tilde{X}$.

Let $T > 0$, $K \in \mathbb{N}, K \geq 2$, $\tau = \frac{T}{K}$. Assume that $\tau R < 1$.

Let $\overline{G} \in B(\{0,1,...,K\};\tilde{X})$ be such that $G^0 = 0$ and consider problem (5.3) with $\varphi^k = \mathcal{R}G^k$ for $k = 1,...,K$ and $U^0 = 0$. Then for $\overline{U} \in B(\{0,1,...,K\};X)$ which solves (5.3) one has

$$\|\mathscr{E}\overline{U}^k\|_{2+2\theta} \leq c\Big(\max_{0\leq k\leq K}\|G^k\|_{2\theta} + \max_{0\leq k_1<k_2\leq K}((k_2 - k_1)\tau)^{-\theta}p(G^{k_2} - G^{k_1})\Big), \tag{5.15}$$

for $k = 0,1,...,K$, where c is a positive constant depending only on θ, R, ϕ_0, M, T and independent of τ_n and $\overline{G}$.

Now we consider the Crank-Nicolson scheme: we replace (5.3) with (5.5). Theorem 5.16 has the following correspondent:

Theorem 5.17 *[91] Assume that the assumptions of Theorem 5.16 are all satisfied and, moreover,*

(f) $\|\tau A\|_{B(X)} \leq S$ with some $S > 0$;

(g) if $|\lambda| \geq 2S$, then

$$\|\mathscr{E}(\lambda - \tau A)^{-1}\mathcal{R}V\|_{2\theta} \leq M\left(\|V\|_{2\theta} + \tau^{-\theta}p(V)\right) \text{ for every } V \in \tilde{X};$$

(h) $\|\mathscr{E}\mathcal{R}V\|_{2\theta} \leq M\Big(\|V\|_{2\theta} + \tau^{-\theta}p(V)\Big)$ for every $V \in \tilde{X}$;

(i) $2\tau R < 1$.

Let $\overline{G} \in B(\{0,1,...,K\};\tilde{X})$ be such that $G^0 = 0$ and consider problem (5.5) with $\varphi^k = \mathcal{R}G^k$ for $k = 1,...,K$ and $U^0 = 0$. If $U \in B\Big(\{0,1,...,K\};X\Big)$ solves (5.5) for $k = 0,1,...,K$, then

$$\|\mathscr{E}U^k\|_{2+2\theta} \leq c\Big(\max_{0\leq k\leq K}\|G^k\|_{2\theta} + \max_{0\leq k_1<k_2\leq K}((k_2 - k_1)\tau)^{-\theta}p(G^{k_2} - G^{k_1})\Big), \tag{5.16}$$

where c is a positive constant depending only on $\theta, R, \phi_0, M, S, T$ and is independent of τ_n and $\overline{G}$.

An application of Theorems 5.16 and 5.17 to the discretization of problem (5.14) is the following. Let $K, n \in I\!N$. We put $\tau := \frac{T}{K}, h := \frac{L}{n}$. We assume that $K \geq 2, n \geq 3$. For $j = 0, 1, ..., n$ we set $a_j := a(jh)$, $b_j = \frac{1}{2}b(jh), c_j := c(jh), \mathbf{N}_{n-1} := \{1, ..., n-1\}$, $\overline{\mathbf{N}}_n := \{0, 1, ..., n-1, n\}$,

$$X := B(\mathbf{N}_{n-1}), \tilde{X} := B(\overline{\mathbf{N}}_n). \tag{5.17}$$

If $V \in X$, we put, as before for $i \in \mathbf{N}_{n-1}$,

$$(A_h V)_i := a_i \frac{\tilde{V}_{i+1} - 2\tilde{V}_i + \tilde{V}_{i+1}}{h^2} + b_i \frac{\tilde{V}_{i+1} - \tilde{V}_{i-1}}{2h} + c_i \tilde{V}_i, \tag{5.18}$$

where

$$\tilde{V}_i = (\mathscr{E}V)_i = \begin{cases} V_i & \text{if } 1 \leq i \leq n-1, \\ 0 & \text{if } i \in \{0, n\}. \end{cases}$$

Next, we define

$$\mathcal{R} \in B(\tilde{X}, X), \mathcal{R}V := V|_{\mathbf{N}_{n-1}} \tag{5.19}$$

for every $V \in \tilde{X}$. Then, again for $V \in \tilde{X}$, we set, for $\theta \in (0, \frac{1}{2})$,

$$\|V\|_{2\theta} := \max\{\|V\|_{\tilde{X}}, \max_{0 \leq i_1 < i_2 \leq n}((i_2 - i_1)h)^{-2\theta}|V_{i2} - V_{i_1}|\}, \tag{5.20}$$

$$\|V\|_{2+2\theta} := \max\{\|V\|_{\tilde{X}}, \max_{0 \leq i \leq n-1}|(\partial_h V)_i|, \max_{0 \leq i \leq n-2}|(\partial_h^2 V)_i|,$$

$$\max_{0 \leq i_1 < i_2 \leq n-2}((i_2 - i_1)h)^{-2\theta}|(\partial_h^2 V)_{i2} - (\partial_h^2 V)_{i_1}|\}, \tag{5.21}$$

with

$$(\partial_h V)_i := \frac{V_{i+1} - V_i}{h} \text{ for } 0 \leq i \leq n-1, (\partial_h^2 V)_i := \frac{V_{i+2} - 2V_{i+1} + V_i}{h} \text{ for } 0 \leq i \leq n-2,$$

$$p(V) := \max\{|V_0|, |V_n|\}. \tag{5.22}$$

One has the following result:

Theorem 5.18 *[91] With the positions (5.17),(5.18), (5.19) and (5.20) the assumptions $(a) - (e)$ of Theorem 5.16 are satisfied, with R, ϕ_0, M independent of h. If we impose the further condition*

$$\tau_n \leq \alpha h^2, \tag{5.23}$$

the same holds true also for assumptions $(f) - (h)$ of Theorem 5.17 (with even S independent of n).

As a consequence, we have:

Theorem 5.19 *[92] Consider the problem (5.14) under the assumptions of Theorem 5.15. With the conventions (5.17),(5.18), (5.19), (5.20), set $G_j^k := f(k\tau_n, jh)$ for $k \in \{1, ..., K\}, j = 0, ..., n$. Put $\varphi^k := \mathcal{R}G^k$ and indicate with G^0 the zero in $B(\overline{\mathbf{N}}_n)$.*

Then if τ_n is sufficiently small,the problem

$$\begin{array}{l} \frac{\tilde{U}_j^k - \tilde{U}_j^{k-1}}{\tau_n} = a_i \frac{\tilde{U}_{i+1}^k - 2\tilde{U}_i^k + \tilde{U}_{i-1}^k}{h^2} + b_i \frac{\tilde{U}_{i+1}^k - \tilde{U}_{i-1}^k}{2h} + c_i \tilde{U}_i^k + \varphi_j^k, \ , \\ \tilde{U}_j^0 = 0, \end{array} \tag{5.24}$$

for $j \in 1, \ldots, n-1$, $k \in \{1, ..., K\}$ has a unique solution such that

$$\|\tilde{U}^k\|_{C_h^{2+2\theta}(\overline{\mathbf{N}}_n)} \leq c\Big(\|f\|_{B([0,T];C^{2\theta}([0,L]))} + \max\{\|f(\cdot, 0)\|_{C^\theta([0,T])}, \|f(\cdot, L)\|_{C^\theta([0,T])}\}\Big), \tag{5.25}$$

with c independent of h and τ_n.

An analogous result holds true for the Crank-Nicholson scheme (5.5). Then we set $G_j^k := f((k-\frac{1}{2})\tau_n, jh)$, under the further condition (5.23).

Remark 5.4 *From Theorem 5.18 it follows that, for the scheme (5.5) with $u_n^0 = 0$ and under condition (5.23)*

$$\|A_h U\|_{C_h^{2\theta}(\mathbf{N}_{n-1})} \le c\Big(\|f\|_{B([0,T];C^{2\theta}([0,L]))} + \max\{\|f_n(\cdot,0)\|_{C^\theta([0,T])}, \|f(\cdot,L)\|_{C^\theta([0,T])}\}\Big), \tag{5.26}$$

with c independent of h and τ_n. In the quoted papers Theorems 5.16 and 5.17 are applied also to the discretization of the heat equation in a square.

A counter example in [91] and [90] shows that condition (5.23) cannot be removed in general. Finally estimates of order of convergence are given in [90].

6 Approximations of Semilinear Equations

Let us consider in a Banach space E the semilinear Cauchy problem

$$u'(t) = Au(t) + f(t, u(t)), \;\; u(0) = u^0, \tag{6.1}$$

with operator A, which generates analytic C_0-semigroup of type $\omega(A) < 0$ and function f is smooth enough. Existence and uniqueness of solution of problem (6.1) have been discussed in [9], [28], [93], [95], [136] for example.

6.1 Approximations of Cauchy problem

By a semidiscrete approximation of the problem (6.1) we mean the Cauchy problems in the Banach spaces E_n

$$u_n'(t) = A_n u_n(t) + f_n(t, u_n(t)), \;\; u_n(0) = u_n^0, \tag{6.2}$$

where the operators A_n generate analytic semigroups in E_n, A_n and A are compatible, functions f_n approximate f and $u_n^0 \to u^0$.

Let Ω be an open set in a Banach space F and let $\mathcal{B} : \bar{\Omega} \to F$ be a compact operator having no fixed points on the boundary of Ω. Then for the vector field $\mathcal{F}(x) = x - \mathcal{B}x$ there is defined the rotation $\gamma(I - \mathcal{B}; \partial\Omega)$, being an integer-valued characteristic of this field. Let x^* be a unique isolated fixed point of the operator $\mathcal{B}$ in the ball S_{r_0} of radius r_0 with the centre at x^*. Then $\gamma(I - \mathcal{B}; \partial S_r) = \gamma(I - \mathcal{B}; \partial S_{r_0})$ for $0 < r \le r_0$, and this common value of the rotations is called the index of the fixed point x^* and it is denoted by *ind* x^*.

Theorem 6.1 *[163] Assume that conditions (A) and (B_1) hold and compact resolvents $(\lambda I - A)^{-1}, (\lambda I - A_n)^{-1}$ converge $(\lambda I - A_n)^{-1} \to (\lambda I - A)^{-1}$ compactly for some $\lambda \in \rho(A)$ and $u_n^0 \to u^0$. Assume that*

(i) the functions f_n, f are bounded and sufficiently smooth so that there exists a unique mild solution $u^*(\cdot)$ of the problem (6.1) on $[0,T]$ (in this situation ind $u^*(\cdot) = 1$);

(ii) $f_n(t, x_n) \to f(t, x)$ uniformly with respect to $t \in [0,T]$ for $x_n \to x$;

(iii) the space E is separable .

Then for almost all n the problems (6.2) have in the neighbourhood of $p_n u^*(\cdot)$ mild solutions $u_n^*(t), t \in [0,T]$. Each sequence $\{u_n^*(t)\}$ is $\mathcal{P}$-compact and $u_n^*(t) \to u^*(t)$ uniformly with respect to $t \in [0,T]$.

Let us consider time discretization with respect to the explicit difference scheme

$$\frac{U_n(t+\tau_n)-U_n(t)}{\tau_n} = A_nU_n(t)+f_n(t,U_n(t)),\ U_n(0)=u_n^0,\ t=k\tau_n, k=\{0,\cdots,K\}. \tag{6.3}$$

Theorem 6.2 *[163] Assume that the conditions of Theorem 6.1 and condition (3.6) are satisfied. Then the functions $U_n(t)$ from (6.3) give an approximate mild solution $u^*(\cdot)$ of the problem (6.1) and, moreover, $U_n(t)\to u^*(t)$ uniformly with respect to $t\in[0,T]$.*

Let us define the operator $\Re(u_n)(t)=u_n(t)-\int_0^t \exp((t-s)A_n)f(s,u_n(s))ds$.

Remark 6.1 *If we assume that the conditions of Theorem 6.1 hold and the functions $f(\cdot), f_n(\cdot)$ have Fréchet derivatives in some balls containing the solutions u^* and u_n^* and moreover, assume that $f'_{n_u}(t,p_nu^*(t))$ are uniformly continuous with respect to the first and second arguments and $f'_{n_{u_n}}(t,u_n(t))\to f'_u(t,u^*(t))$ uniformly with respect to $t\in[0,T]$ for $u_n\to u^*$, then [163] for almost all n the problems (6.2) have in the neighbourhood of $p_nu^*(\cdot)$ mild solutions $u_n^*(t), t\in[0,T]$. Each sequence $\{u_n^*(\cdot)\}$ is $\mathcal{P}$-compact and $u_n^*(t)\to u^*(t)$ uniformly with respect to $t\in[0,T]$ and, moreover, for sufficiently large $n\ge n_0$ and some $T^*\le T$ we have*

$$c_1\ \epsilon_n(u^*,u_n^0)\le \|u_n^*-p_nu^*\|_{F_n}\le c_2\ \epsilon_n(u^*,u_n^0),$$

where the constants c_1,c_2 are not dependent on n, $F_n=C([0,T];E_n)$, while

$$\epsilon_n(u^*,u_n^0)=\max_{t\in[0,T^*]}\|\Re(p_nu^*)(t)-\exp(tA_n)u_n^0\|_{E_n}.$$

Let $\mathcal{U}_n(t)=(I_n+\tau_nA_n)^k, \Im_n(u_n)(t)=u_n(t)-\Sigma_{l=1}^{k-1}\mathcal{U}_n((k-l)\tau_n)\ f_n(l\tau_n,u_n(l\tau_n))\ \tau_n$.

Remark 6.2 *If we assume that the conditions of Theorem 6.2 hold and the functions $f(\cdot), f_n(\cdot)$ have Fréchet derivatives in some balls containing the solutions $u^*(\cdot)$ and $u_n^*(\cdot)$ and moreover, assume that $f'_{n_{u_n}}(t,p_nu^*(t))$ are uniformly continuous with respect to the first and second arguments and $f'_{n_{u_n}}(t,u_n(t))\to f'_u(t,u^*(t))$ uniformly with respect to $t\in[0,T]$ for $u_n\to u^*$, and condition (3.6) holds, then [163] the functions $U_n(t)$ from (6.3) give an approximate mild solution of the problem (6.1) and $U_n^*(t)\to u^*(t)$ uniformly with respect to $t\in[0,T]$ and , moreover, for sufficiently large $n\ge n_0$ and some $T^*\le T$ we have*

$$c_1\ \epsilon_n(u^*,u_n^0)\le \|U_n^*-p_nu^*\|_{F_n^{\tau_n}}\le c_2\ \epsilon_n(u^*,u_n^0),$$

where the constants c_1,c_2 do not depend on n, $F_n^{\tau_n}=\{u_n(k\tau_n):\max_{0\le k\tau_n\le T}\ \|u_n(k\tau_n)\|_{E_n}\ <\infty\}$, while $\epsilon_n(u^*,u_n^0)=\max_{t\in[0,T^*]}\|\Im_n(p_nu^*)(t)-\mathcal{U}_n(t)u_n^0\|_{E_n}$.

Schemes which have higher order of convergence than (6.3) are considered in [143], [163]. The Runge-Kutta methods for semilinear equations were considered in [76], [134], [133], [132], [143], [146].

6.2 Approximation of periodic problem

Let us consider in a Banach space E the semilinear T-periodic problem

$$v'(t)=Av(t)+f(t,v(t)),\ \ v(t)=v(T+t), t\in\overline{IR}_+, \tag{6.4}$$

with operator A, which generates an analytic compact C_0-semigroup and with the function f which is smooth enough and $f(t,x)=f(t+T,x)$ for any $x\in E$ and $t\in\overline{IR}_+$. Let $u(\cdot;u^0)$ be the solution of

Cauchy problem (6.1) with initial data $u(0;u^0)=u^0$. This function $u(\cdot;u^0)$ is also a mild solution, i.e. it satisfies the integral equation

$$u(t)=\exp(tA)u^0+\int_0^t \exp\left((t-s)A\right)f(s,u(s))ds,\ t\in \overline{I\!R}_+. \tag{6.5}$$

Then the shift operator $\mathcal{K}(u^0)=u(T;u^0)$ can be defined and it maps E to E. If $u(\cdot;x^*)$ is a periodic solution of (6.1), then x^* is a zero of the compact vector field defined by $I-\mathcal{K}$, i.e. $\mathcal{K}(x^*)=x^*$.

Remark 6.3 *We assume here that operator $(I-\exp(TA))^{-1}$ exists and is bounded. In the mean time it is just enough to assume that $(I-\exp(tA))^{-1}\in B(E)$ holds for $t\geq t_0$ with some $t_0>0$. This assumption is not restrictive as without loss of generality we are able to change A to $A-\omega I$ and get $\|\exp\left(t(A-\omega)\right)\|\leq Me^{-\delta t}$ for $\delta>0, t\geq 0$. It follows then [26] that $(I-\exp(tA))^{-1}\in B(E)$ for any $t>0$.*

Remark 6.4 *We said that function f is smooth enough in the sense that it has to be at least continuous in both arguments, $\sup_{t\in[0,T],\|x\|\leq c_1}\|f(t,x)\|\leq C_2$ and such that there exists the global mild solution of the problem $u'(t)=Au(t)+f(t,u(t)),\ \ u(0)=u^0, t\in \overline{I\!R}_+$.*

Definition 6.1 *The solution $u(\cdot)$ of Cauchy problem (6.1) is said to be stable in the Lyapunov sense if for any $\epsilon>0$ there is $\delta>0$ such that inequality $\|u(0)-\tilde{u}(0)\|\leq\delta$ implies $\max_{0\leq t<\infty}\|u(t)-\tilde{u}(t)\|\leq\epsilon$, where $\tilde{u}(\cdot)$ is a mild solution of (6.1) with the initial value $\tilde{u}(0)$.*

Definition 6.2 *The solution $u(\cdot)$ of Cauchy problem (6.1) is said to be uniformly asymptotically stable at the point $u(0)$ if it is stable in the Lyapunov sense and for any mild solution $\tilde{u}(\cdot)$ of (6.1) with $\|u(0)-\tilde{u}(0)\|\leq\delta$ it follows that $\lim_{t\to\infty}\|u(t)-\tilde{u}(t)\|=0$ uniformly in $\tilde{u}(\cdot)\in B(u(0);\delta)$, i.e. there is a function $\phi_{u(0),\delta}(\cdot)$ such that $\|u(t;u(0))-u(t;\tilde{u}(0))\|\leq\phi_{u(0),\delta}(t)$ with $\phi_{u(0),\delta}(t)\to 0$ as $t\to\infty$ and $\|u(0)-\tilde{u}(0)\|\leq\delta$.*

Constructive conditions on operator A and f to ensure that equation $u'(t)=Au(t)+f(u(t)), u(u)=u^0$ is asymptotically $k-$dimensional are given in [170], [169]. They concern the location of eigenvalues of A, i.e. $\lambda_{k+1}-\lambda_k>2L, \lambda_{k+1}>L$.

Theorem 6.3 *[35] Assume that conditions (A) and (B") hold and compact resolvents $(\lambda I-A)^{-1}, (\lambda I-A_n)^{-1}$ converge $(\lambda I-A_n)^{-1}\to(\lambda I-A)^{-1}$ compactly for some $\lambda\in\rho(A)$. Assume that*

(i) the functions f, f_n are sufficiently smooth, so that there exists an isolated mild solution $v^*(\cdot)$ of periodic problem (6.4) with $v^*(0)=x^*$ such that the Cauchy problem (6.1) with $u(0)=x^*$ has an uniformly asymptotically stable isolated solution at the point x^* (in this case $ind\, v^*(\cdot)=1$);

(ii) $f_n(t,x_n)\to f(t,x)$ for any $t\in[0,T]$ as $x_n\to x$;

(iii) the space E is separable .

Then for almost all n the problems

$$v_n'(t)=A_nv_n(t)+f_n(t,v_n(t)), v_n(t)=v_n(t+T), t\in\overline{I\!R}_+, \tag{6.6}$$

have periodic mild solutions $v_n^*(t), t\in[0,T]$, in the neighbourhood of $p_nv^*(\cdot)$, where $v^*(\cdot)$ is a mild periodic solution of (6.4) with $v^*(0)=x^*$. Each sequence $\{v_n^*(\cdot)\}$ is $\mathcal{P}$-compact and $v_n^*(t)\to v^*(t)$ uniformly with respect to $t\in[0,T]$.

We say that a fixed point x^* to operator $\mathcal{K}$ in Banach lattice E is stable from above [95] if given $\epsilon>0$, there is a $\delta>0$ such that $\|\mathcal{K}^kx-x^*\|\leq\epsilon$ for all $k\in I\!N$ if $x^*\preceq x$ and $\|x-x^*\|\leq\delta$. Using this notion Theorem 6.3 can be reformulated for positive semigroups due to result from [59].

Theorem 6.4 *Let the operators A_n and A from the problems (6.4) and (6.6) be consistent and let E, E_n be order unit spaces and $e_n \in D(A_n) \cap intE_n^+$. Assume that the operators A_n have the POD property and $A_n e_n \preceq 0$ for sufficiently large n and compact resolvents $(\lambda I - A)^{-1}, (\lambda I - A_n)^{-1}$ converge $(\lambda I - A_n)^{-1} \to (\lambda I - A)^{-1}$ compactly for some $\lambda \in \rho(A)$. Assume that*

(i) the functions f, f_n are sufficiently smooth, bounded and positive, so that there exists a mild solution $u^*(\cdot)$ of the Cauchy problem (6.3) such that elements $u^*(0) = x^*$ is a stable from above and fixed points of operator $\mathcal{K}$, with $x^* \prec y, \mathcal{K}y \preceq y$ (in this situation ind $x^* = 1$);

(ii) $f_n(t, x_n) \to f(t, x)$ uniformly with respect to $t \in [0, T]$ for $x_n \to x$;

(iii) the space E is separable .

Then for almost all n the problems (6.6) have periodic mild solutions $v_n^*(t), t \in [0, T]$ in the neighbourhood of $p_n v^*(\cdot)$, where $v^*(\cdot)$ is any stable from above mild periodic solution of (6.4). Each sequence $\{v_n^*(\cdot)\}$ is $\mathcal{P}$-compact and $v_n^*(t) \to v^*(t)$ uniformly with respect to $t \in [0, T]$.

Remark 6.5 *The technique which is used here could be applied to the case of condensing operators [3]. For example, the resolvent of Δ in $L^2(\mathbb{R}^d)$ is condensing, but it is not compact.*

In [117] they study the qualitative behavior of spatially semidiscrete finite element solutions of a semilinear parabolic problem near an unstable hyperbolic equilibrium.

The shadowing approach to study the long-time behavior of numerical approximations of semilinear parabolic equations was studied in [116].

ACKNOWLEDGEMENTS

The authors acknowledge the support of NATO-CP Advanced Fellowship Programme of TÜBITAK (Turkish Scientific Research Council), University of Antwerpen, Russian Foundation for Basic Research (01-01-00398) and University of Bologna. We would also like to thank Professor E.H. Twizell for his valuable comments.

References

[1] N. H. Abdelaziz. On approximation by discrete semigroups. *J. Approx. Theory*, 73(3):253–269, 1993.

[2] N. H. Abdelaziz and Paul R. Chernoff. Continuous and discrete semigroup approximations with applications to the Cauchy problem. *J. Operator Theory*, 32(2):331–352, 1994.

[3] S. B. Agase and V. Raghavendra. Existence of mild solutions of semilinear differential equations in Banach spaces. *Indian J. Pure Appl. Math.*, 21(9):813–821, 1990.

[4] M. Ahués. A class of strongly stable operator approximations. *J. Austral. Math. Soc. Ser. B*, 28(4):435–442, 1987.

[5] M. Ahués and F. Hocine. A note on spectral approximation of linear operations. *Appl. Math. Lett.*, 7(2):63–66, 1994.

[6] M. Ahués and A. Largillier. Two numerical approximations for a class of weakly singular integral operators. *Appl. Numer. Math.*, 17(4):347–362, 1995.

[7] M. Ahues and S. Piskarev. Spectral approximation of weakly singular integral operators. i: Convergence theory. In et. al. Constanda. C., editor, *Integral methods in science and engineering. Vol. II: Approximation methods. Proceedings of the 4th international conference, IMSE '96, Oulu, Finland, June 17–20*, Harlow, 1996. Longman, Pitman Res.

[8] H. A. Alibekov and P. E. Sobolevskii. Stability and convergence of difference schemes of a high order of approximation for parabolic partial differential equations. *Ukrain. Mat. Zh.*, 32(3):291–300, 1980.

[9] H. Amann. *Linear and quasilinear parabolic problems. Vol. I.* Birkhäuser Verlag, Basel, 1995. Abstract linear theory.

[10] P. M. Anselone. *Collectively compact operator approximation theory and applications to integral equations.* Prentice-Hall Inc., Englewood Cliffs, N. J., 1971. With an appendix by Joel Davis, Prentice-Hall Series in Automatic Computation.

[11] P. M. Anselone. Nonlinear operator approximation. In *Moderne Methoden der numerischen Mathematik (Tagung 200-Jahrfeier, Tech. Univ. Clausthal, Clausthal, 1975)*, volume 32 of *Internat. Ser. Numer. Math.*, pages 17–24. Birkhäuser, Basel, 1976.

[12] P. M. Anselone. Recent advances in operator approximation theory. In *Methods of functional analysis in approximation theory (Bombay, 1985)*, volume 76 of *Internat. Schriftenreihe Numer. Math.*, pages 309–324. Birkhäuser, Basel, 1986.

[13] P. M. Anselone and T. W. Palmer. Spectral properties of collectively compact sets of linear operators. *J. Math. Mech.*, 17:853–860, 1967/1968.

[14] P. M. Anselone and T. W. Palmer. Collectively compact sets of linear operators. *Pacific J. Math.*, 25:417–422, 1968.

[15] P. M. Anselone and M. L. Treuden. Regular operator approximation theory. *Pacific J. Math.*, 120(2):257–268, 1985.

[16] P. M. Anselone and M. L. Treuden. Spectral analysis of asymptotically compact operator sequences. *Numer. Funct. Anal. Optim.*, 17(7-8):679–690, 1996.

[17] F. Aràndiga and V. Caselles. Approximations of positive operators and continuity of the spectral radius. III. *J. Austral. Math. Soc. Ser. A*, 57(3):330–340, 1994.

[18] F. Aràndiga and V. Caselles. On strongly stable approximations. *Rev. Mat. Univ. Complut. Madrid*, 7(2):207–217, 1994.

[19] W. Arendt. Vector-valued Laplace transforms and Cauchy problems. *Israel J. Math.*, 59(3):327–352, 1987.

[20] A. Ashyralyev. Difference schemes of higher order accuracy for elliptic equations. In *Application of the methods of functional analysis to nonclassical equations in mathematical physics (Russian)*, pages 3–14. Akad. Nauk SSSR Sibirsk. Otdel. Inst. Mat., Novosibirsk, 1989.

[21] A. Ashyralyev, S. Piskarev, and L. Weis. On well-posedness of difference schemes for abstract parabolic equations in $L^p([0,T];E)$ spaces. *Submitted.*

[22] A. Ashyralyev and P. E. Sobolevskiĭ. *Well-posedness of parabolic difference equations*, volume 69 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1994. Translated from the Russian by A. Iacob.

[23] A. O. Ashyralyev. Some difference schemes for parabolic equations with nonsmooth initial data. *Izv. Akad. Nauk Turkmen. SSR Ser. Fiz.-Tekhn. Khim. Geol. Nauk*, 6:70–72, 1983.

[24] N. Yu. Bakaev, S. Larsson, and V. Thomée. Backward Euler type methods for parabolic integro-differential equations in Banach space. *RAIRO Modél. Math. Anal. Numér.*, 32(1):85–99, 1998.

[25] M. E. Ballotti, J. A. Goldstein, and M. E. Parrott. Almost periodic solutions of evolution equations. *J. Math. Anal. Appl.*, 138(2):522–536, 1989.

[26] V. Barbu and N. H. Pavel. On the invertibility of $I \pm \exp(-tA)$, $t > 0$, with A maximal monotone. In *World Congress of Nonlinear Analysts '92, Vol. I–IV (Tampa, FL, 1992)*, pages 2231–2237. de Gruyter, Berlin, 1996.

[27] B. Baumer and F. Neubrander. Existence and uniqueness of solutions of ordinary linear differential equations in Banach spaces. Preprint, Departmeent of Mathematics, 1996. Louisiana state Univ., Baton Rouge.

[28] Ph. Benilan, N. Crandall, and A. Pazy. Nonlinear evolution equations in Banach spaces. book in print, 1998.

[29] J. Bergh and J. Löfström. *Interpolation spaces. An introduction.* Springer-Verlag, Berlin, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.

[30] A. Bobrowski. Degenerate convergence of semigroups. *Semigroup Forum*, 49(3):303–327, 1994.

[31] A. Bobrowski. Integrated semigroups and the trotter-kato theorem. *Bull. Polish Acad. Sci.*, 41(4):297–303, 1994.

[32] A. Bobrowski. Examples of a pointwise convergence of semigroups. *Ann. Univ. Mariae Curie-Skłodowska Sect. A*, 49:15–33, 1995.

[33] A. Bobrowski. Generalized telegraph equation and the Sova-Kurtz version of the Trotter-Kato theorem. *Ann. Polon. Math.*, 64(1):37–45, 1996.

[34] A. Bobrowski. On the generation of non-continuous semigroups. *Semigroup Forum*, 54(2):237–252, 1997.

[35] N. A. Bobylev, J. K. Kim, S. K. Korovin, and S. Piskarev. Semidiscrete approximations of semilinear periodic problems in Banach spaces. *Nonlinear Anal.*, 33(5):473–482, 1998.

[36] P. Bolley, J. Camus, and Phạm T. L. Estimation de la résolvante du problème de Dirichlet dans les espaces de Hölder. *C. R. Acad. Sci. Paris Sér. I Math.*, 305(6):253–256, 1987.

[37] R. Bouldin. Operator approximations with stable eigenvalues. *J. Austral. Math. Soc. Ser. A*, 49(2):250–257, 1990.

[38] J. H. Bramble, J. E. Pasciak, P. H. Sammon, and V. Thomée. Incomplete iterations in multistep backward difference methods for parabolic problems with smooth and nonsmooth data. *Math. Comp.*, 52(186):339–367, 1989.

[39] P. Brenner, M. Crouzeix, and V. Thomée. Single-step methods for inhomogeneous linear differential equations in Banach space. *RAIRO Anal. Numer.*, 16:5–26, 1982.

[40] P. Brenner, V. Thomée, and L. B. Wahlbin. *Besov spaces and applications to difference methods for initial value problems.* Springer-Verlag, Berlin, 1975. Lecture Notes in Mathematics, Vol. 434.

[41] P. Brenner and Thomée V. On rational approximations of semigroups. *SIAM J. Numer. Anal.*, 16:683–694, 1979.

[42] K. Burrage and S. Piskarev. Stochastic methods for ill-posed problems. *BIT*, 40(2):226–240, 2000.

[43] S. Busenberg and B. Wu. Convergence theorems for integrated semigroups. *Differential Integral Equations*, 5(3):509–520, 1992.

[44] P. L. Butzer, W. Dickmeis, L. Hahn, and R. J. Nessel. Lax-type theorems and a unified approach to some limit theorems in probability theory with rates. *Resultate Math.*, 2:30–53, 1979.

[45] P. L. Butzer, W. Dickmeis, Hu. Jansen, and R. J. Nessel. Alternative forms with orders of the Lax equivalence theorem in Banach spaces. *Computing*, 17(4):335–342, 1976/77.

[46] P. L. Butzer, W. Dickmeis, and R. J. Nessel. Lax-type theorems with orders in connection with inhomogeneous evolution equations in Banach spaces. In *Linear spaces and approximation (Proc. Conf., Math. Res. Inst., Oberwolfach, 1977)*, volume 21 of *Lecture Notes in Biomath.*, pages 531–546. Springer, Berlin, 1978.

[47] P. L. Butzer, W. Dickmeis, and R. J. Nessel. Numerical analysis of evolution equations. In *Symposium on Numerical Analysis, Kyoto, July 11–12, 1978*, volume 1 of *Lecture Notes in Numerical and Applied Analysis*, pages 1–163. Kinokuniya Book Store Co., Ltd., Tokyo, 1979.

[48] A. S. Carasso, J. G. Sanderson, and J. M. Hyman. Digital removal of random media image degradations by solving the diffusion equation backwards in time. *SIAM J. Numer. Anal.*, 15(2):344–367, 1978.

[49] F. Chatelin. *Spectral approximation of linear operators.* Computer Science and Applied Mathematics. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1983. With a foreword by P. Henrici, With solutions to exercises by Mario Ahués.

[50] A.J. Chorin, M.F. McCracken, T. J. R. Hughes, and J. E. Marsden. Product formulas and numerical algorithms. *Comm. Pure Appl. Math.*, 31(2):205–256, 1978.

[51] M. Crouzeix. On multistep approximation of semigroups in Banach spaces. In *Proceedings of the 2nd international conference on computational and applied mathematics (Leuven, 1986)*, volume 20, pages 25–35, 1987.

[52] M. Crouzeix, S. Larsson, S. Piskarev, and V. Thomée. The stability of rational approximations of analytic semigroups. *BIT*, 33(1):74–84, 1993.

[53] M. Crouzeix, S. Larsson, and V. Thomée. Resolvent estimates for elliptic finite element operators in one dimension. *Math. Comp.*, 63(207):121–140, 1994.

[54] M. Crouzeix and V. Thomée. On the discretization in time of semilinear parabolic equations with nonsmooth initial data. *Math. Comp.*, 49(180):359–377, 1987.

[55] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.

[56] Yu. L. Dalecky and S. V. Fomin. *Measures and differential equations in infinite-dimensional space*, volume 76 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1991. Translated from the Russian, With additional material by V. R. Steblovskaya, Yu. V. Bogdansky [Yu. V. Bogdanskiĭ] and N. Yu. Goncharuk.

[57] Yu. L. Dalecky and N. Yu. Goncharuk. Random operator functions defined by stochstic differential equations. In *Measures and differential equations in infinite-dimensional space*. 1990. Translated from the Russian. With additional material by V. R. Steblovskaya, Yu. V. Bogdansky.

[58] Yu-L. Dalecky and N-Yu. Goncharuk. On a quasilinear stochastic differential equation of parabolic type. *Stochastic Anal. Appl.*, 12(1):103–129, 1994.

[59] E. N. Dancer. Upper and lower stability and index theory for positive mappings and applications. *Nonlinear Anal.*, 17:205–217, 1991.

[60] E. B. Davies and M. M. H. Pang. The Cauchy problem and a generalization of the Hille- Yosida theorem. *Proc. London Math. Soc. (3)*, 55(1):181–208, 1987.

[61] R. deLaubenfels. Integrated semigroups, C-semigroups and the abstract Cauchy problem. *Semigroup Forum*, 41(1):83–95, 1990.

[62] R. deLaubenfels. C-semigroups and the Cauchy problem. *J. Funct. Anal.*, 111(1):44–61, 1993.

[63] R. J. Dickmeis, W.and Nessel. On uniform boundedness principles and Banach-Steinhaus theorems with rates. *Numer. Funct. Anal. Optim.*, 3:19–52, 1981.

[64] W. Dickmeis and R. J. Nessel. Classical approximation processes in connection with Lax equivalence theorems with orders. *Acta Sci. Math. (Szeged)*, 40(1-2):33–48, 1978.

[65] W. Dickmeis and R. J. Nessel. A unified approach to certain counterexamples in approximation theory in connection with a uniform boundedness principle with rates. *J. Approx. Theory*, 31:161–174, 1981.

[66] B. Eberhardt and G. Greiner. Baillon's theorem on maximal regularity. *Acta Appl. Math.*, 27:47–54, 1992. Positive operators and semigroups on Banach lattices (Curaçao, 1990).

[67] S. N. Ethier and T. G. Kurtz. *Markov processes*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1986. Characterization and convergence.

[68] H. Fujita. On the finite element approximation for parabolic equations: an operator theoretical approach. In *Computing methods in applied sciences and engineering (Second Internat. Sympos., Versailles, 1975), Part 1*, volume 134 of *Lecture Notes in Econom. and Math. Systems*, pages 171–192. Springer, Berlin, 1976.

[69] H. Fujita and A. Mizutani. On the finite element method for parabolic equations. I. Approximation of holomorphic semi-groups. *J. Math. Soc. Japan*, 28(4):749–771, 1976.

[70] H. Fujita and T. Suzuki. On the finite element approximation for evolution equations of parabolic type. In *Computing methods in applied sciences and engineering (Proc. Third Internat. Sympos., Versailles, 1977), I*, volume 704 of *Lecture Notes in Math.*, pages 207–221. Springer, Berlin, 1979.

[71] H. Fujita and T. Suzuki. Evolution problems. In *Handbook of numerical analysis, Vol. II*, pages 789–928. North-Holland, Amsterdam, 1991.

[72] J. A. Goldstein. *Semigroups of linear operators and applications.* Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 1985.

[73] J. A. Goldstein. A survey of semigroups of linear operators and applications. In *Semigroups of linear and nonlinear operations and applications (Curaçao, 1992)*, pages 9–57. Kluwer Acad. Publ., Dordrecht, 1993.

[74] J. A. Goldstein and Jr. Sandefur, J. T. An abstract d'Alembert formula. *SIAM J. Math. Anal.*, 18(3):842–856, 1987.

[75] J.A. Goldstein, R. deLaubenfels, and Jr. Sandefur, J. T. Regularized semigroups, iterated Cauchy problems and equipartition of energy. *Monatsh. Math.*, 115(1-2):47–66, 1993.

[76] C. González and C. Palencia. Stability of Runge-Kutta methods for abstract time-dependent parabolic problems: the Hölder case. *Math. Comp.*, 68:73–89, 1999.

[77] E. Görlich and D. Pontzen. An approximation theorem for semigroups of growth order α and an extension of the trotter-lie formula. *J. Funct. Anal.*, 50(3):414–425, 1983.

[78] R. D. Grigorieff. Zur Theorie linearer approximationsregulärer Operatoren. I, II. *Math. Nachr.*, 55:233–249; ibid. 55 (1973), 251–263, 1973.

[79] R. D. Grigorieff. Diskrete Approximation von Eigenwertproblemen. I. Qualitative Konvergenz. *Numer. Math.*, 24(4):355–374, 1975.

[80] R. D. Grigorieff. Diskrete Approximation von Eigenwertproblemen. II. Konvergenzordnung. *Numer. Math.*, 24(5):415–433, 1975.

[81] R. D. Grigorieff. Über diskrete Approximationen nichtlinearer Gleichungen 1. Art. *Math. Nachr.*, 69:253–272, 1975.

[82] R. D. Grigorieff. Diskrete Approximation von Eigenwertproblemen. III. Asymptotische Entwicklungen. *Numer. Math.*, 25(1):79–97, 1975/76.

[83] R. D. Grigorieff. Zur Charakterisierung linearer approximationsregulärer Operatoren. In *Mathematical papers given on the occasion of Ernst Mohr's 75th birthday*, pages 63–77. Tech. Univ. Berlin, Berlin, 1985.

[84] R. D. Grigorieff. Some stability inequalities for compact finite difference schemes. *Math. Nachr.*, 135:93–101, 1988.

[85] R. D. Grigorieff. Time discretization of semigroups by the variable two-step BDF method. In *Numerical treatment of differential equations (Halle, 1989)*, volume 121 of *Teubner-Texte Math.*, pages 204–216. Teubner, Stuttgart, 1991.

[86] R. D. Grigorieff and H. Jeggle. Approximation von Eigenwertproblemen bei nichlinearer Parameterabhängigkeit. *Manuscripta Math.*, 10:245–271, 1973.

[87] D. Guidetti. Optimal regularity for mixed parabolic problems in spaces of functions which are Hölder continuous with respect to space variables. Technical report, Department of Mathematics, University of Bologna, 1995.

[88] D. Guidetti. The parabolic mixed Cauchy-Dirichlet problem in spaces of functions which are Hölder continuous with respect to space variables. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 7(3):161–168, 1996.

[89] D. Guidetti. The parabolic mixed Cauchy-Dirichlet problem in spaces of functions which are Hölder continuous with respect to space variables. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 7(3):161–168, 1996.

[90] D. Guidetti and S. Piskarev. On maximal regularity of difference schemes for parabolic equations in $c^{\theta}(\overline{\omega})$ spaces. N 8, Dipartimento di Matematica, Universitaá degli Studi di Bologna, 1997. Preprint.

[91] D. Guidetti and S. Piskarev. On maximal regularity of Crank-Nicholson schemes for parabolic equations in $C^{\theta}(\overline{\Omega})$ space. In *Proccedings of International Conference "Function spaces. Differential operators. Problems in mathematical education*, volume 2, pages 2231–2237. RUDN, Moscow, 1998.

[92] D. Guidetti and S. Piskarev. Stability of Rothe's scheme and maximal regularity for parabolic equations in $C^{\theta}(\overline{\Omega})$. In *Progress in partial differential equations, Vol. 1 (Pont-à-Mousson, 1997)*, pages 167–180. Longman, Harlow, 1998.

[93] D. Henry. *Geometric theory of semilinear parabolic equations.* Springer-Verlag, Berlin, 1981.

[94] R. Hersh and T. Kato. High-accuracy stable difference schemes for well-posed initial value problems. *SIAM J. Numer. Anal.*, 16(4):670–682, 1979.

[95] P. Hess. *Periodic-parabolic boundary value problems and positivity.* Longman Scientific & Technical, Harlow, 1991.

[96] R. H. W. Hoppe. A constructive approach to the Bellman semigroup. *Nonlinear Anal.*, 9(11):1165–1181, 1985.

[97] V. K. Ivanov, V. V. Vasin, and V. P. Tanana. *Theory of linear ill-posed problems and its applications.* Nauka, Moscow, 1978.

[98] N. Kalton and L. Weis. The H^{∞} calculus and sums of closed operators. *submitted.*

[99] N. Kalton and L. Weis. Perturbation and interpolation theorems for the H^{∞} calculus. *in preparation.*

[100] N.J. Kalton and G. Lancien. A solution to the problem of L^p-maximal regularity. Math. Z., 235(3):559–568, 2000.

[101] S. Kanda. Convergence of difference approximations and nonlinear semigroups. *Proc. Amer. Math. Soc.*, 108(3):741–748, 1990.

[102] T. Kato. *Perturbation theory for linear operators.* Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.

[103] N. Kenmochi and S. Oharu. Difference approximation of nonlinear evolution equations and semi-groups of nonlinear operators. *Publ. Res. Inst. Math. Sci.*, 10(1):147–207, 1974/75.

[104] J. Kisyński. A proof of the Trotter-Kato theorem on approximation of semi-groups. *Colloq. Math.*, 18:181–184, 1967.

[105] Y. Kobayashi. Difference approximation of Cauchy problems for quasi-dissipative operators and generation of nonlinear semigroups. *J. Math. Soc. Japan*, 27(4):640–665, 1975.

[106] Y. Kobayashi. Difference approximation of evolution equations and generation of nonlinear semi-groups. *Proc. Japan Acad.*, 51(6):406–410, 1975.

[107] S. G. Krein. *Linear differential equations in Banach space.* American Mathematical Society, Providence, R.I., 1971. Translated from the Russian by J. M. Danskin, Translations of Mathematical Monographs, Vol. 29.

[108] P. Kunstmann and L. Weis. Perturbation theorems for maximal regularity. *submitted.*

[109] P.Ch. Kunstmann Abstrakte Cauchy probleme und Distributionshalbgruppen. Dissertation, Univ. Kiel, 1995.

[110] P.Ch. Kunstmann. Regularization of semigroups that are strongly continuous for $t > 0$. Proc. Amer. Math. Soc., 1998, 126, N9, p. 2721-2724.

[111] T. G. Kurtz. Extensions of Trotter's operator semigroup approximation theorems. *J. Functional Analysis*, 3:354–375, 1969.

[112] T. G. Kurtz. A general theorem on the convergence of operator semigroups. *Trans. Amer. Math. Soc.*, 148:23–32, 1970.

[113] T. G. Kurtz and P. E. Protter. Weak convergence of stochastic integrals and differential equations. II. Infinite-dimensional case. In *Probabilistic models for nonlinear partial differential equations (Montecatini Terme, 1995)*, volume 1627 of *Lecture Notes in Math.*, pages 197–285. Springer, Berlin, 1996.

[114] R. Labbas. Some results on the sum of linear operators with nondense domains. *Ann. Mat. Pura Appl. (4)*, 154:91–97, 1989.

[115] A. Largillier. A numerical quadrature for some weakly singular integral operators. *Appl. Math. Lett.*, 8(1):11–14, 1995.

[116] S. Larsson and S. Yu. Pilyugin. Numerical shadowing near the global attractor for a semilinear parabolic equation. Preprint 21, Department of Mathematics, Chalmers University of Technology, 1998.

[117] S. Larsson and J. M. Sanz-Serna. The behavior of finite element solutions of semilinear parabolic problems near stationary points. *SIAM J. Numer. Anal.*, 31(4):1000–1018, 1994.

[118] S. Larsson and J.-M. Sanz-Serna. A shadowing result with applications to finite element approximation of reaction-diffusion equations. *Math. Comp.*, 68(225):55–72, 1999.

[119] S. Larsson, V. Thomée, and L. B. Wahlbin. Numerical solution of parabolic integro-differential equations by the discontinuous Galerkin method. *Math. Comp.*, 67(221):45–71, 1998.

[120] S. Larsson, V. Thomée, and S. Z. Zhou. On multigrid methods for parabolic problems. *J. Comput. Math.*, 13(3):193–205, 1995.

[121] I. Lasiecka and A. Manitius. Differentiability and convergence rates of approximating semigroups for retarded functional-differential equations. *SIAM J. Numer. Anal.*, 25(4):883–907, 1988.

[122] P. D. Lax and R. D. Richtmyer. Survey of the stability of linear finite difference equations. *Comm. Pure Appl. Math.*, 9:267–293, 1956.

[123] R. D. Lazarov, V. L. Makarov, and A. A. Samarskii. Application of exact difference schemes for constructing and investigating difference schemes on generalized solutions. *Mat. Sb. (N.S.)*, 117(159)(4):469–480, 559, 1982.

[124] C. Le Merdy. Counterexamples on L_p-maximal regularity. *Math. Z.*, 230:47–62, 1999.

[125] M. Le Roux and V. Thomée. Numerical solution of semilinear integrodifferential equations of parabolic type with nonsmooth data. *SIAM J. Numer. Anal.*, 26(6):1291–1309, 1989.

[126] T. D. Lee. Difference equations and conservation laws. *J. Statist. Phys.*, 46(5-6):843–860, 1987.

[127] H. Linden. Starke Konvergenz im verallgemeinerten Sinne und Spektra. *Math. Z.*, 134:205–213, 1973.

[128] H. Linden. Über die Stabilität von Eigenwerten. *Math. Ann.*, 203:215–220, 1973.

[129] C. Lizama. On an extension of the Trotter-Kato theorem for resolvent families of operators. *J. Integral Equations Appl.*, 2(2):269–280, 1990.

[130] C. Lizama. On the convergence and approximation of integrated semigroups. *J. Math. Anal. Appl.*, 181(1):89–103, 1994.

[131] G. J. Lord and A. M. Stuart. Discrete Gevrey regularity, attractors and upper-semicontinuity for a finite difference approximation to the Ginzburg-Landau equation. *Numer. Funct. Anal. Optim.*, 16(7-8):1003–1047, 1995.

[132] C. Lubich and A. Ostermann. Runge-Kutta methods for parabolic equations and convolution quadrature. *Math. Comp.*, 60(201):105–131, 1993.

[133] C. Lubich and A. Ostermann. Runge-Kutta approximation of quasi-linear parabolic equations. *Math. Comp.*, 64(210):601–627, 1995.

[134] C. Lubich and A. Ostermann. Runge-Kutta time discretization of reaction-diffusion and Navier-Stokes equations: nonsmooth-data error estimates and applications to long-time behaviour. *Appl. Numer. Math.*, 22(1-3):279–292, 1996. Special issue celebrating the centenary of Runge-Kutta methods.

[135] Ch. Lubich, I. H. Sloan, and V. Thomée. Nonsmooth data error estimates for approximations of an evolution equation with a positive-type memory term. *Math. Comp.*, 65(213):1–17, 1996.

[136] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*, volume 16 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Basel, 1995.

[137] I. V. Mel'nikova and S. V. Bochkareva. C-semigroups and the regularization of an ill-posed Cauchy problem. *Dokl. Akad. Nauk*, 329(3):270–273, 1993.

[138] M. Miklavčič. Approximations for weakly nonlinear evolution equations. *Math. Comp.*, 53(188):471–484, 1989.

[139] Jr. Mills, W. H. The resolvent stability condition for spectra convergence with application to the finite element approximation of noncompact operators. *SIAM J. Numer. Anal.*, 16:695–703, 1979.

[140] I. Miyadera and N. Tanaka. Exponentially bounded C-semigroups and generation of semigroups. *J. Math. Anal. Appl.*, 143(2):358–378, 1989.

[141] J. K. Mountain. The lax equivalence theorem for linear, inhomogeneous equations in l^2 spaces. *J. Approx. Theory*, 33:126–130, 1981.

[142] M. T. Nair. On strongly stable approximations. *J. Austral. Math. Soc. Ser. A*, 52(2):251–260, 1992.

[143] E. Nakaguchi and A. Yagi. Error estimate of implicit runge-kutta methods for quasilinear abstract equations of parabolic type in banach space. Preprint, Department of Applied Physics, Graduate School of Engineering, Osaka University, 1996. apanese Journal of Mathematics.

[144] O. Nevanlinna. On the convergence of difference approximations to nonlinear contraction semigroups in Hilbert spaces. *Math. Comp.*, 32(142):321–334, 1978.

[145] O. Nevanlinna and G. Vainikko. Limit spectrum of discretely converging operators. *Numer. Funct. Anal. Optim.*, 17(7-8):797–808, 1996.

[146] A. Ostermann and M. Roche. Runge-Kutta methods for partial differential equations and fractional orders of convergence. *Math. Comp.*, 59(200):403–420, 1992.

[147] C. Palencia. A stability result for sectorial operators in Banach spaces. *SIAM J. Numer. Anal.*, 30(5):1373–1384, 1993.

[148] C. Palencia. On the stability of variable stepsize rational approximations of holomorphic semigroups. *Math. Comp.*, 62(205):93–103, 1994.

[149] C. Palencia. Stability of rational multistep approximations of holomorphic semigroups. *Math. Comp.*, 64(210):591–599, 1995.

[150] C. Palencia. Maximum norm analysis of completely discrete finite element methods for parabolic problems. *SIAM J. Numer. Anal.*, 33(4):1654–1668, 1996.

[151] C. Palencia and B. García-Archilla. Stability of linear multistep methods for sectorial operators in Banach spaces. *Appl. Numer. Math.*, 12(6):503–520, 1993.

[152] C. Palencia and S. Piskarev. On multiplicative perturbations of c_0-groups and c_0-cosine operator functions. Technical report, Department of Mathematics, Universidad de Valladolid, Valladolid, Spain, 1998.

[153] C. Palencia and J. M. Sanz-Serna. Equivalence theorems for incomplete spaces: an appraisal. *IMA J. Numer. Anal.*, 4(1):109–115, 1984.

[154] C. Palencia and J. M. Sanz-Serna. An extension of the Lax-Richtmyer theory. *Numer. Math.*, 44(2):279–283, 1984.

[155] A. K. Pani, V. Thomée, and L. B. Wahlbin. Numerical methods for hyperbolic and parabolic integro-differential equations. *J. Integral Equations Appl.*, 4(4):533–584, 1992.

[156] S. V. Parter. On the roles of "stability" and "convergence" in semidiscrete projection methods for initial-value problems. *Math. Comp.*, 34(149):127–154, 1980.

[157] D. Pfeifer. Some general probabilistic estimations for the rate of convergence in operator semigroup representations. *Appl. Anal.*, 23(1-2):111–118, 1986.

[158] S. Piskarev. Approximation of holomorphic semigroups. *Tartu Riikl. Ül. Toimetised*, (492):3–14, 1979.

[159] S. Piskarev. Error estimates in the approximation of semigroups of operators by Padé fractions. *Izv. Vyssh. Uchebn. Zaved. Mat.*, (4):33–38, 1979.

[160] S. Piskarev. Stability of difference schemes in Cauchy problems with almost periodic solutions. *Differentsial' nye Uravneniya*, 20(4):689–695, 1984.

[161] S. Piskarev. Principles of discretization methods III. Report ak -3410, Acoustic Institute, Academy of Scince USSR, 1986. 87 pages.

[162] S. Piskarev. Estimates for the rate of convergence in the solution of ill-posed problems for evolution equations. *Izv. Akad. Nauk SSSR Ser. Mat.*, 51(3):676–687, 1987.

[163] S. Piskarev. Convergence of difference schemes in the solution of nonlinear parabolic equations. *Mat. Zametki*, 44(1):112–123, 156, 1988.

[164] S. Piskarev. A method of increasing the acuuracy of the solution of initial value problems in Banach space. *USSR Comput. Math. Phys.*, 28(6):211–212, 1988.

[165] S. Piskarev. Approximation of positive C_0-semigroups of operators. *Differentsial' nye Uravneniya*, 27(7):1245–1250, 1287, 1991.

[166] S. Piskarev. On the approximation of perturbed C_0-semigroups. *Differentsial'nye Uravneniya*, 30(2):339–341, 367, 1994.

[167] R. D. Richtmyer and K. W. Morton. *Difference methods for initial-value problems.* Interscience Publishers John Wiley & Sons, Inc., New York-London-Sydney, 1967. Second edition. Interscience Tracts in Pure and Applied Mathematics, No. 4.

[168] A. V. Romanov. Conditions for asymptotic k-dimensionality of semilinear parabolic equations. *Uspekhi Mat. Nauk*, 46(1(277)):213–214, 1991.

[169] A. V. Romanov. Sharp estimates for the dimension of inertial manifolds for nonlinear parabolic equations. *Izv. Ross. Akad. Nauk Ser. Mat.*, 57(4):36–54, 1993.

[170] A. V. Romanov. On the limit dynamics of evolution equations. *Uspekhi Mat. Nauk*, 51(2(308)):173–174, 1996.

[171] E. B. Saff, A. Schönhage, and R. S. Varga. Geometric convergence to e^{-z} by rational functions with real poles. *Numer. Math.*, 25(3):307–322, 1975/76.

[172] E. B. Saff and R. S. Varga. On the zeros and poles of Padé approximants to e^z. *Numer. Math.*, 25(1):1–14, 1975/76.

[173] E. B. Saff and R. S. Varga. Geometric overconvergence of rational functions in unbounded domains. *Pacific J. Math.*, 62(2):523–549, 1976.

[174] E. B. Saff and R. S. Varga. On the zeros and poles of Padé approximants to e^z. II. In *Padé and rational approximation (Proc. Internat. Sympos., Univ. South Florida, Tampa, Fla., 1976)*, pages 195–213. Academic Press, New York, 1977.

[175] E. B. Saff and R. S. Varga. On the zeros and poles of Padé approximants to e^z. III. *Numer. Math.*, 30(3):241–266, 1978.

[176] A. Samarskii and V. Andréev. *Méthodes aux différences pour équations elliptiques.* Éditions Mir, Moscow, 1978. Traduit du russe par Djilali Embarek.

[177] A. Samarskii and A.V. Gulin. *Stability of difference schemes.* Nauka, Moscow, 1973. (in Russian).

[178] J. M. Sanz-Serna and C. Palencia. A general equivalence theorem in the theory of discretization methods. *Math. Comp.*, 45(171):143–152, 1985.

[179] S.-Y. Shaw and Ch.-Ch. Kuo. Local C-semigrous and the Cauchy problems. Preprint, Departmeent of Mathematics, 1996. National Central University, Taiwan.

[180] P. E. Sobolevskii. The coercive solvability of difference equations. *Dokl. Akad. Nauk SSSR*, 201:1063–1066, 1971.

[181] P. E. Sobolevskii. The theory of semigroups and the stability of difference schemes. In *Operator theory in function spaces (Proc. School, Novosibirsk, 1975) (Russian)*, pages 304–337, 344. Izdat. "Nauka" Sibirsk. Otdel., Novosibirsk, 1977.

[182] P. E. Sobolevskii and Hoáng Văn Lai. Difference schemes of optimal type for the approximate solution of parabolic equations (the Banach case). *Ukrain. Mat. Zh.*, 33:39–46, 1981.

[183] Z. Strkalj and L. Weis. On operator-valued fourier multiplier theorems. *submitted.*

[184] F. Stummel. Diskrete Konvergenz linearer Operatoren. III. In *Linear operators and approximation (Proc. Conf., Oberwolfach, 1971)*, pages 196–216. Internat. Ser. Numer. Math., Vol. 20. Birkhäuser, Basel, 1972.

[185] T. Takahashi. Convergence of difference approximation of nonlinear evolution equations and generation of semigroups. *J. Math. Soc. Japan*, 28(1):96–113, 1976.

[186] N. Tanaka. Approximation of integrated semigroups by "integrated" discrete parameter semigroups. *Semigroup Forum*, 55(1):57–67, 1997.

[187] N. Tanaka and I. Miyadera. Some remarks on C-semigroups and integrated semigroups. *Proc. Japan Acad. Ser. A Math. Sci.*, 63(5):139–142, 1987.

[188] N. Tanaka and N. Okazawa. Local C-semigroups and local integrated semigroups. *Proc. London Math. Soc. (3)*, 61(1):63–90, 1990.

[189] V. Thomée. Finite difference methods for linear parabolic equations. In *Handbook of numerical analysis, Vol. I*, pages 5–196. North-Holland, Amsterdam, 1990.

[190] V. Thomée. Approximate solution of O.D.E.s in Banach space—rational approximation of semigroups. In *Hellenic research in mathematics and informatics '92 (Athens, 1992)*, pages 409–420. Hellenic Math. Soc., Athens, 1992.

[191] V. Thomée. Finite element methods for parabolic problems—some steps in the evolution. In *Finite element methods (Jyväskylä, 1993)*, volume 164 of *Lecture Notes in Pure and Appl. Math.*, pages 433–442. Dekker, New York, 1994.

[192] A. N. Tihonov and V. Ja. Arsenin. *Methods for the solution of ill-posed problems.* Nauka, Moscow, 1979.

[193] A. N. Tikhonov and V. Y. Arsenin. *Solutions of ill-posed problems.* V. H. Winston & Sons, Washington, D.C.: John Wiley & Sons, New York, 1977. Translated from the Russian, Preface by translation editor Fritz John, Scripta Series in Mathematics.

[194] H. F. Trotter. Approximation of semi-groups of operators. *Pacific J. Math.*, 8:887–919, 1958.

[195] H. F. Trotter. On the product of semi-groups of operators. *Proc. Amer. Math. Soc.*, 10:545–551, 1959.

[196] H. F. Trotter. Approximation and perturbation of semigroups. In *Linear operators and approximation, II (Proc. Conf., Oberwolfach Math. Res. Inst., Oberwolfach, 1974)*, pages 3–21. Internat. Ser. Numer. Math., Vol. 25. Birkhäuser, Basel, 1974.

[197] E. H. Twizell. *Computational methods for partial differential equations.* Ellis Horwood Ltd., Chichester, 1984.

[198] E. H. Twizell and A. Q. M. Khaliq. Multiderivative methods for periodic initial value problems. *SIAM J. Numer. Anal.*, 21(1):111–122, 1984.

[199] T. Ushijima. On the abstract Cauchy problems and semi-groups of linear operators in locally convex spaces. *Sci. Papers College Gen. Ed. Univ. Tokyo*, 21:93–122, 1971.

[200] T. Ushijima. Approximation theory for semi-groups of linear operators and its application to approximation of wave equations. *Japan. J. Math. (N.S.)*, 1(1):185–224, 1975/76.

[201] T. Ushijima. The approximation of semi-groups of linear operators and the finite element method. In *Boundary value problems for linear evolution: partial differential equations (Proc. NATO Advanced Study Inst., Liège, 1976)*, volume 29 of *NATO Advanced Study Inst. Ser., Ser. C: Math. and Phys. Sci.*, pages 351–362. Reidel, Dordrecht, 1977.

[202] T. Ushijima. Approximation of semigroups and the finite element method. *Sûgaku*, 32:133–148, 1980.

[203] T. Ushijima and M. Matsuki. Fully discrete approximation of a second order linear evolution equation related to the water wave problem. In *Functional analysis and related topics, 1991 (Kyoto)*, volume 1540 of *Lecture Notes in Math.*, pages 361–380. Springer, Berlin, 1993.

[204] G. Vainikko. *Funktionalanalysis der Diskretisierungsmethoden.* B. G. Teubner Verlag, Leipzig, 1976. Mit Englischen und Russischen Zusammenfassungen, Teubner-Texte zur Mathematik.

[205] G. Vainikko. Über die Konvergenz und Divergenz von Näherungsmethoden bei Eigenwertproblemen. *Math. Nachr.*, 78:145–164, 1977.

[206] G. Vainikko. Über Konvergenzbegriffe für lineare Operatoren in der numerischen Mathematik. *Math. Nachr.*, 78:165–183, 1977.

[207] G. Vainikko. Approximative methods for nonlinear equations (two approaches to the convergence problem). *Nonlinear Anal.*, 2:647–687, 1978.

[208] G. Vainikko. Foundations of finite difference method for eigenvalue problems. In *The use of finite element method and finite difference method in geophysics (Proc. Summer School, Liblice, 1977)*, pages 173–192. Česk. Akad. Věd, Prague, 1978.

[209] G. Vainikko and O. Karma. An estimate for the rate of convergence of eigenvalues in approximation methods. *Tartu Riikl. Ül. Toimetised*, (833):75–83, 1988.

[210] G. Vainikko and S. Piskarev. Regularly compatible operators. *Izv. Vysš. Učebn. Zaved. Matematika*, 10(185):25–36, 1977.

[211] J. van Casteren, S. Piskarev, and S.-Y. Shaw. Discretisation of C-semigroups. submitted to J. of Inverse and Ill-posed Problem.

[212] J. van Casteren, S. Piskarev, and S.-Y. Shaw. Discretization of C-semigroups. Technical report, Proceedings of the Conference on inverse and ill-posed problems, Moscow State University, 1998. 18.

[213] V. V. Vasilev, S. G. Krein, and S. Piskarev. Operator semigroups, cosine operator functions, and linear differential equations. In *Mathematical analysis, Vol. 28 (Russian)*, Itogi Nauki i Tekhniki, pages 87–202, 204. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1990. Translated in J. Soviet Math. **54** (1991), no. 4, 1042–1129.

[214] V.V. Vasiliev and S. Piskarev. Diffeential equations in Banach spaces II. Cosine-operator functions. Translated in J. Soviet Math.

[215] V.V. Vasiliev and S. Piskarev. Diffeential equations in Banach spaces I. Semigroup theory. Moscow State University Publishing House, 1996. in Russian, 164 pages.

[216] V. V. Vasilev, S.I. Piskarev. Bibliography on Differential Equations in Abstract Spaces. Moscow State University, 2001, Electronic journal.

(http://www.srcc.msu.su/num_anal/list_wrk/page_5u.htm)

PostScript (1115,2 Kb) PostScript.zip (336,1 Kb)

[217] J. Voigt. On the convex compactness property for the strong operator topology. *Note Mat.*, 12:259–269, 1992. Dedicated to the memory of Professor Gottfried Köthe.

[218] L. Weis. A new approach to maximal L_p regularity. *Proc of the 6th Internat. Conf on Evolution Equations*, Marcel Dekker 2000.

[219] L. Weis. Operator-valued multiplier theorems and maximal L_p regularity. *Mathematische Annalen*, to appear.

[220] R. Wolf. Über lineare approximationsreguläre Operatoren. *Math. Nachr.*, 59:325–341, 1974.

[221] L. Wu. The semigroup stability of the difference approximations for initial-boundary value problems. *Math. Comp.*, 64(209):71–88, 1995.

[222] Q. Zheng and Y. S. Lei. Exponentially bounded C-semigroup and integrated semigroup with nondensely defined generators. I. Approximation. *Acta Math. Sci. (English Ed.)*, 13(3):251–260, 1993.

Davide Guidetti
Department of Mathematics, University of Bologna
Piazza di Porta S. Donato, 5, 40127 Bologna, Italy
guidetti@dm.unibo.it

Bülent Karasözen
Department of Mathematics, Middle East Technical University
06531 Ankara, Turkey
bulent@metu.edu.tr

Serguei Piskarev
Scientific Research Computer Center, Moscow State University
Vorobjevy Gory, Moscow 119899, Russia
serguei@piskarev.srcc.msu.su